Review Article

Javad Sharifi-Rad*, Cristina Quispe, Aknur Turgumbayeva, Zehra Mertdinç, Sena Tütüncü, Elif Feyza Aydar, Beraat Özçelik, Stępień-Warda Anna, Staniak Mariola, Anna Koziróg, Anna Otlewska, Hubert Antolak, Surjit Sen, Krishnendu Acharya, Natallia Lapava, Simin Emamzadeh-Yazdi, Miquel Martorell, Manoj Kumar, Elena Maria Varoni, Marcello Iriti* and Daniela Calina*

Santalum Genus: phytochemical constituents, biological activities and health promoting-effects

https://doi.org/10.1515/znc-2022-0076 Received April 2, 2022; accepted July 15, 2022; published online September 7, 2022

Abstract: *Santalum* genus belongs to the family of Santalaceae, widespread in India, Australia, Hawaii, Sri Lanka, and Indonesia, and valued as traditional medicine,

Medicina, Universidad del Azuay, Cuenca, Ecuador, E-mail: javad.sharifirad@gmail.com. https://orcid.org/0000-0002-7301-8151 (J. Sharifi-Rad); **Marcello Iriti**, Department of Biomedical, Surgical and Dental Sciences, Università degli Studi di Milano, 20133 Milano, Italy; and National Interuniversity Consortium of Materials Science and Technology (INSTM), 50121 Firenze, Italy, E-mail: marcello.iriti@unimi.it. https://orcid.org/ 0000-0002-5063-1236 (M. Iriti); and **Daniela Calina**, Department of Clinical Pharmacy, University of Medicine and Pharmacy of Craiova, 200349 Craiova, Romania,

E-mail: calinadaniela@gmail.com

Cristina Quispe, Facultad de Ciencias de la Salud, Universidad Arturo Prat, Avda. Arturo Prat 2120, 1110939, Iquique, Chile, E-mail: elquispe@unap.cl

Aknur Turgumbayeva, Higher School of Medicine, Al-Farabi Kazakh National University, Almaty, Kazakhstan; and School of Pharmacy, JSC "S. D. Asfendiyarov Kazakh National Medical University", Almaty, Kazakhstan, E-mail: aknurturgumbayeva@gmail.com

Zehra Mertdinç, Sena Tütüncü and Elif Feyza Aydar, Faculty of Chemical and Metallurgical Engineering, Department of Food Engineering, Istanbul Technical University, 34469 Maslak, Istanbul, Turkey, E-mail: mertdincz@itu.edu.tr (Z. Mertdinç),

tutuncus@itu.edu.tr (S. Tütüncü), aydar@itu.edu.tr (E.F. Aydar) Beraat Özçelik, Faculty of Chemical and Metallurgical Engineering, Department of Food Engineering, Istanbul Technical University, 34469 Maslak, Istanbul, Turkey; and BIOACTIVE Research & Innovation Food Manufacturing Industry Trade LTD Co., Maslak, Istanbul 34469, Turkey, E-mail: ozcelik@itu.edu.tr

Stępień-Warda Anna and Staniak Mariola, Department of Forage Crop Production, Institute of Soil Science and Plant Cultivation – State Research Institute, Czartoryskich 8, 24-100 Puławy, Poland, E-mail: astepien@iung.pulawy.pl (S.-W. Anna), staniakm@iung.pulawy.pl (S. Mariola) rituals and modern bioactivities. Sandalwood is reported to possess a plethora of bioactive compounds such as essential oil and its components (α -santalol and β -santalol), phenolic compounds and fatty acids. These bioactives play important role in contributing towards biological activities and health-promoting effects in humans. Pre-clinical and clinical studies have shown the role of sandalwood

Anna Koziróg, Anna Otlewska and Hubert Antolak, Faculty of Biotechnology and Food Sciences, Lodz University of Technology, Institute of Fermentation Technology and Microbiology, Wolczanska 171/173, 90 – 924 Lodz, Poland, E-mail: anna.kozirog@p.lodz.pl (A. Koziróg), anna.otlewska@p.lodz.pl (A. Otlewska), hubert.antolak@p.lodz.pl (H. Antolak)

Surjit Sen, Molecular and Applied Mycology and Plant Pathology Laboratory, Department of Botany, University of Calcutta, 700019, Kolkata, India; and Department of Botany, Fakir Chand College, Diamond Harbour, West Bengal, 743331, India,

E-mail: surjitsen09@gmail.com. https://orcid.org/0000-0002-4038-7292

Krishnendu Acharya, Department of Botany, Fakir Chand College, Diamond Harbour, West Bengal, 743331, India,

E-mail: krish_paper@yahoo.com **Natallia Lapava,** Medicine Standartization Department of Vitebsk State Medical University, Vitebsk, Republic of Belarus, E-mail: natallia_karazhan@tut.by

Simin Emamzadeh-Yazdi, Department of Plant and Soil Sciences, University of Pretoria, Gauteng 0002, Pretoria, South Africa, E-mail: simin.emamzadeh.yazdi@gmail.com

Miquel Martorell, Department of Nutrition and Dietetics, Faculty of Pharmacy, Centre for Healthy Living, University of Concepción, 4070386 Concepción, Chile; and Universidad de Concepción, Unidad de Desarrollo Tecnológico, UDT, 4070386 Concepción, Chile, E-mail: martorellpons@gmail.com

Manoj Kumar, Chemical and Biochemical Processing Division, ICAR – Central Institute for Research on Cotton Technology, 400019 Mumbai, India, E-mail: manojkumarpuniya114@gmail.com

Elena Maria Varoni, Department of Biomedical, Surgical and Dental Sciences, Università degli Studi di Milano, 20133 Milano, Italy; and National Interuniversity Consortium of Materials Science and Technology (INSTM), 50121 Firenze, Italy, E-mail: elena.varoni@unimi.it

Open Access. © 2022 the author(s), published by De Gruyter. 📷 This work is licensed under the Creative Commons Attribution 4.0 International License.

^{*}Corresponding authors: Javad Sharifi-Rad, Facultad de

extract as antioxidant, anti-inflammatory, antibacterial, antifungal, antiviral, neuroleptic, antihyperglycemic, antihyperlipidemic, and anticancer activities. Safety studies on sandalwood essential oil (EO) and its extracts have proven them as a safe ingredient to be utilized in health promotion. Phytoconstituents, bioactivities and traditional uses established sandalwood as one of the innovative materials for application in the pharma, food, and biomedical industry.

Keywords: bioactivities; clinical studies; essential oil; phytochemistry; safety; sandalwood.

1 Introduction

The genus *Santalum* is a woody flowering plant that belongs to the family Santalaceae commonly known as sandalwood. The members of the genus are generally trees or shrubs. The plant is obligate hemiparasite attaching itself by haustoria to establish contact with the host and extracts xylem sap for nutrients and water [1]. The family Santalaceae comprises 29 genera with around 400 species out of which 18 well-recognized species are under the genus *Santalum* [1–7] (Table 1).

Sandalwood is generally popular for its fragrant heartwood oil used by cosmetic industries for the production of perfume [13, 18–21]. The high demand for sandalwood oil and timber has resulted in drastic overharvesting; as a result, many taxa are now considered as rare, threatened or listed as endangered [22]. Moreover, one species *Santalum fernandezianum* Phil. from the Juan Fernandez Islands (South Pacific Ocean), has been reported extinct due to over-exploitation by human beings [23]. About 25 species belong to the genus *Santalum*, they are evergreen trees or shrubs characterized by a semi-parasitic lifestyle. They conduct photosynthesis; however, they take in water and inorganic nutrients by parasitizing on the roots of other plant species.

Plants of the genus *Santalum* are characterized by the production of EOs with many biological properties due to the high content of bioactive substances such as lignans, glycosides, triterpenoids, and sesquiterpenoids (α/β -santalol - the compound found in the largest amount). These bioactive compounds include antioxidant, anti-inflammatory, antibacterial, antifungal, antiviral, neuroleptic, antihyperglycemic, antihyperlipidemic, and anticancer activities.

Santalum genus has been known to possess many health benefits proved based on traditional uses and modern biological approaches through preclinical studies. Traditionally, *Santalum* genus has been used as an antipyretic, immune booster, antidiarrhea, and for treating cold and cough. Modern uses have shown their effect as antioxidant, anti-inflammatory, antibacterial,

Taxonomic group/Section	Species and IUCN Red List Category	Common name	Geographical Occurrence	Reference
Section <i>Santalum</i> skottsb.	S. album L.	Indian sandalwood	Australia, Belgium, Cambodia, China, Germany, Great britain, Holand, India, Indonesia, Japan, Madagaskar, Malaysia, Norway, Spain, Srilanka, Switzerland, and the United States.	[3, 8, 9]
Section <i>Solenantha</i> Tuyama	S. fernandezianum Phil.	Freycinet sandalwood, or iliahi	Hawaiian islands (Oʻahu, Molokaʻi)	[10, 11]
	S. haleakalae Hillebr.	Haleakala sandalwood or iliahi	Hawaiian islands (Maui)	[11, 12]
	S. pyrularium A. Gray	Hawaiian sandalwood or iliahi	Hawaiian islands (Kauaʻi)	[3, 5, 13, 14]
Section <i>Hawaiiensia</i> skottsb.	S. ellipticum Gaudich.	Coastal sandalwood or 'Iliahialo'e	Hawaiian islands	[3, 11, 12]
	S. paniculatum Hook. & Arn.	Hawaiʻi	Hawaiian islands	[3, 11]
Section <i>Polynesica</i> skottsb.	S. fernandezianum F.Phil.	Chile sandalwood	Juan fernandez islands	[3]
Genus <i>Eucarya</i> T.Mitch.	S. acuminatum (R.Br.) A.DC.	Desert Quandong, native Peach	Australia	[15–17]

Table 1: Section, recognized species, according to the International Union for Conservation of Nature (IUCN) category, common name and geographical distribution of *Santalum* species.

antifungal, antiviral, neuroleptic, antihyperglycemic, antihyperlipidemic, and anticancer activities. One of the most important parameters considered for the application of plant extracts in the biomedical field is its safety. The safety aspects of EO components have been studied by various researchers and it is concluded that extracts from *Santalum* genus are fairly safe to be used as healthpromoting effects. The current review is the first of its kind that gives a snapshot of the *Santalum* genus concerning its traditional uses, bioactive components, bioactivities (*in vitro*, *in vivo* and clinical trials), and its safety aspects while using it as a health-promoting agent in humans.

2 Review methodology

Available information on the genus *Santalum*, its biological properties, and its potential mechanisms of action was collected by searching the following databases: PubMed/ Medline, Web of Science and ScienceDirect. The following MeSH terms were used: "Santalum/growth & development", "Santalum/chemistry", "Plant Oils/isolation & purification", "Animals", "Apoptosis/drug effects", "Carcinogenesis/drug effects", "Cell Cycle Checkpoints/drug effects", "Humans", "Mice", "Plant Extracts/chemistry", "Plant Extracts/therapeutic use", "Plant Oils/therapeutic use", "Antioxidants/ metabolism".

The study included research articles and reviews published in extenso, written in English language in scientific journals, book chapters and books with information about *Santalum* genum and sandalwood. Editorials/letters to publishers, case reports, conference abstracts, studies that included homeopathic preparations were excluded. The PlantList database was used to verify the taxonomy and provide information on the classification and distribution of *Santalum* subspecies [24, 25].

3 Botany

Santalum is widely distributed to semi-arid areas from Indonesia in the West to Juan Fernandez Islands (Chile) in the East and from Hawaiian Archipelago in the North to New Zealand in the South [8] (Figure 1). The major production places of the plant are shown in Table 1.

The well-recognized species are broadly grouped into four categories viz. Indian sandalwood (*Santalum album* L.), Australian sandalwood (*Santalum acuminatum* (R. Br.) A. DC.), Hawaiian sandalwood (*Santalum ellipticum* Gaudich., *Santalum freycinetianum* F. Phil., *Santalum haleakalae* Hillebr., *Santalum paniculatum* Hook. & Arn., and *Santalum pyrularium* A. Gray), and Pacific Islands sandalwood (*S. fernandezianum* Phil.).

A taxonomic grouping in *Santalum* is purely based on morphological characters. It has been reported that Section *Santalum* is described as reddish corollas that are longer than wide and partly superior ovaries [12, 26, 27]. Based on smaller ovaries, longer perianth tubes and absence of hairs to the filament [28] separated the two Hawaiian members (*S. freycinetianum* and *S. haleakalae*) from the section



Figure 1: Geographical distribution of Santalum genus.

Santalum into the endemic section *Solenantha*. The characteristic features of the section *Hawaiiensia* are explained as having white, brown, orange or green corollas that are as wide as long and inferior ovaries [12, 26, 27].

Section Polynesica is similar in appearance to the section Hawaiiensia but with partly superior ovaries [26]. A molecular phylogenetic study reveals that the sectional classification of Santalum needs revision [3]. Skottsberg [26, 29] suggested that sections Hawaiiensia and Polynesica were closely related based on morphological characters and section Polynesica was treated as a synonym of section Hawaiiensia by [30]. But molecular phylogenetic analysis indicates that sections Hawaiiensia and Polynesica are not close to one another rather related to other taxa of section Santalum and should not be united taxonomically [3]. Revisionary studies based on molecular data considered six species of Santalum in Hawaii, whereas previously there were only four recognized species [5]. Hawaiian species are considered to be a result of two colonization processes which comprises four species within redflowered section Solenantha (i.e., S. freycinetianum, S. haleakalae, and S. pyrularium); and two species within white-flowered section Hawaiiensia (i.e., S. ellipticum, and S. paniculatum) [14].

General features of the genus *Santalum* are evergreen trees or shrubs; leaves opposite rarely alternate, sometimes in whorls, glabrous or sometimes glaucous, ovate, obovate or lanceolate, coriaceous [31]. Flowers cymose panicle, axillary or in the terminal, tetra or pentamerous, hermaphrodite; bracts small. Perianth-tube campanulate to conical or ovoid, adnate to the base of the ovary; stamens 4-5, dorsifixed, filament slender, short, anthers ovate; 4-lobed; style long, stigma 2-4 lobed; ovary inferior or partly inferior; ovules 2-3. Flowers produce sweet to a week or no fragrance. Fruit globose to sub-globose drupe, annulate on the top by the deciduous perianth; seed subglobose; albumen copious. Morphological differences of some important species are summarized in Table 2.

4 Traditional uses

The close-grained heartwood of *Santalum* is used for ornamental and carving work. *Santalum* fruits are edible and the seeds contain fatty oil which is suitable for the manufacture of paint. Incense sticks are made of powdered heartwood and are used in houses and temples. In addition, powered heartwood is ground into a paste and used as a cosmetic [35]. *Santalum* genus is mentioned in Indian mythology, folklore, scripture, and the oldest literature (for example, Vinaya Pitaka (400–300 BC) and Milinda Pahna (200 BC)) and also in the epic Ramayana and Mahabharata. The ancient Egyptians used *Santalum* plants oil for embalming the dead and in the ritual burning to venerate the gods. In certain communities among the Hindus it is traditional to put a piece of sandalwood in the funeral pyre. A beige-coloured paste obtained from sandalwood is put in on the forehead and other body parts, especially by devotees of God Krishna (Vaishnavites) and for ritual bathing of Hindu gods [36]. In Zoroastrian temples, *Santalum* burns in sacred lights to soothe the problems of all mankind. It is used by Jews, Buddhists, Hindus, as well as almost all other belief systems for its huge variety in attributes [35].

5 Bioactive composition

5.1 Essential oil, terpenes, and derivatives

After 30 years of growth with a natural condition, oil is collected from the heartwood of sandalwood. The yield of the oil depends on the age of the tree; an old mature tree gives an oil yield between 2.5–6%; the colour of the heartwood, individual tree understudy, location within the tree, and the environment of growth of the tree. Sandalwood oil consists of main terpenoids: mono- and sesquiterpenes and their oxygenated derivatives (mostly the alcohols, ketones, and aldehydes) and also some fatty acids, and phenylpropanoids chemical compounds [37–39] (Figure 2).

The bark extract of S. album contains mainly santalol (90%) [40, 41], exo-norbicycloekasantalal, β -santalic, teresantalic, nortricycloekasantalic, bicycloekasantalic, dihydro- β -santalic acids, urs-12-en-3b-il-palmitate, β -sitosterol, (+)epi- β -santalol, (-) β -santalol, (-)trans- β -santalol, α -santalol (52%), β -santalol (23%), epi- β -santalene, cislanceol, cis-nuciferol, β -, epi- β -teresantalic acid, β -, epi- β -norekasantalic acid, β -, epi- β -ekasantalic acid, α -santalic acid, 11-keto-dihydro- α -santalic acid, bisabolenols A, B, C, D and E, tricycloekasan-talol, α - and β -santalenes, trans- α -bergamotene, α -curcumone, nuciferol. The bark extract of S. album includes l-allohydroxiproline, betulinic acid, β -sitosterol, and fatty acids. The bark extract of S. album contains betulinic acid (0.05%), β -sitosterol, glucose, fructose, and sucrose [38, 40]. Although including a low amount of trans- β -santalol, cis-lanceol hydrocarbons, α -santalene, β -santalene, α -bergamotene, epi- β -santalene, as α -curcumene, β -curcumene, γ -curcumene, β -bisabolene and α -bisabolol; cis- α -santalol (53%), cis- β -santalol (23%), α-trans-bergamotol, epi-cis-β-santalol sesquiterpene alcohols are the major components of the sandalwood oil [42-46].

Category	Species	Size	Leaves	Flower	Fruit	Reference
Indian sandalwood	S. album L.	Small to the medium-size tree with slender droop- ing branches	Opposite, lanceolate to ovate; acute to obtuse at base, entire; apex acute to acuminate; pale green to lush green	Initially, straw yellow col- oured and gradually turn to deep purplish or brown	Green to purplish– black; succulent	[8, 9, 31, 32]
Australian sandalwood	S. acuminatum (R. Br.) A. DC.	A shrubby small tree	Opposite, more or less lanceolate; pale green to olive-green; acute apex	Small, creamy white or greenish-white	Globose, green turning to orange– red to bright, glossy red; persistent tepal scar	[15, 16, 33]
Hawaiian sandalwood	S. ellipticum Gaudich.	Shrub to small tree	Elliptic to orbicular, ovate, or obovate; leathery to succulent; glaucous; dull, greyish green	Greenish in bud but tinged with brown, or- ange, or salmon after opening; produce a sweet fragrance; flower as long as wide	Purple to black drupes, with a distinctive apical receptacular ring	[11, 34]
	S. paniculatum Hook. & Arn.	Shrub or tree	Ovate, obovate or elliptic; upper surfaces glossy and lower surface dull; yellowish orange to bluish or olive green.	Greenish in bud but tinged with brown, or- ange, or salmon after opening; produce a sweet fragrance; flowers as long as wide	Purple to black with a distinctive apical receptacular ring.	[11, 34]
	S. freycinetianum F. Phil.	Shrub to tree	Narrowly elliptic, oblong, to narrowly ovate; acute to rounded apex; bit glau- cous; green	Light pink turning deep pink with maturity (rarely with white interiors); pro- duce a weak fragrance; flowers longer than wide	Reddish-purple to almost black with a distinctive sub- apical receptacular ring	[5, 10, 11]
	S. haleakalae Hillebr.	Small tree	Ovate, obovate, or orbic- ular; stiff to coriaceous surfaces; olive green	Deep pink to red throughout, or with white to pink interiors; produce a weak fragrance; flowers longer than wide	Black or purplish- black with a distinc- tive sub-apical receptacular ring.	[5, 11]
	S. pyrularium A. Gray	Small tree or shrubby tree	Opposite; elliptic, ovate, to oblong; glaucous abax- ially not much paler on abaxial surface; acute to obtuse apices; medium to dark green	Cream to purple throughout, greenish with the purple interior, or greenish-white turning red with age	Red, elliptic, with subapical ring	[5]

Table 2: Contrasting morphological characters of important Santalum species.

5.1.1 Extraction of the Santalum Oil

EO is one of the important components of an important component of sandalwood and its isolation from sandalwood depends on the methods of extraction. The EO of the sandalwood is widely used in the fragrance industry due to having a strong aroma and has various biological activities such as: anticancer, antiviral, antidiarrheal, cytotoxic activities, among others [47]. The different extraction methods can be applied to the *Santalum* during oil extraction. Therefore, the composition and amount of the fragrance and volatile compounds found in oil may

vary dependent on the extraction methods. The conventional steam distillation and hydrodistillation methods are performed under high temperatures (around 100 °C) which can often result in loss of volatile compounds and changes in the odour [48–50]. Maceration or Soxhlet type solvent extraction are other techniques that have some the drawbacks such as large volume of solvent usage, exposure to hazardous and flammable liquid organic solvents, and environmental issues [51]. Therefore, the use of some solvent-free "green methods" during the extraction of EO has gained prominence in recent years. Microwave-assisted extraction, subcritical CO_2 (SC– CO_2)



Figure 2: Illustrative scheme with the most important bioactive constituents of Santalum essential oil and their pharmacological properties.

extraction and some other combined novel technologies, such as microwave-assisted hydrodistillation method, are preferred due to having higher selectivity and extraction yield, need for less time for analysis and not posing environmental and safety concerns [52, 53].

Kusuma and Mahfud [52] objected to looking into the effects of the newly employed microwave airhydrodistillation method for extraction of EOs and comparison with classical microwave hydrodistillation method. Results of this research showed that additional airflow to the microwave hydrodistillation can help obtain the sandalwood oil in higher yield directly proportional with air flow rate. The compound composition of microwave air-hydro distilled sandalwood oil is larger than another method concerning identification 43 compounds whereas 37 compounds are recorded in microwave hydrodistillation. Microwave air-hydrodistillation provides better aroma/fragrance quality than microwave hydrodistillation extracts [52].

Nautiyal [54] mentioned that extraction yield and quality affect the trade of sandalwood oil. It was also highlighted that heartwood preparation and the extraction method have an influence on α - and β -santalol levels in the obtained oil. In the study, eight different extraction methods which are SC–CO₂, ethyl alcohol, benzene, diethyl ether, toluene, steam distillation, hydrodistillation, and

alkaline-hydro distillation are examined. The highest yield is obtained from SC-CO₂ extraction, 3.83 grams per liter (g/L). In the analysis of extracted sandalwood oil for α - and β-santalol levels were examined through gas chromatography (GC). The most efficient extraction methods are SC-CO₂, ethyl alcohol, and steam distillation; they include nearly 84% total α - and β - santalol. Hydrodistillation is the least efficient in terms of having α : β - santalol ratio, 3:1, whereas SC-CO₂, ethyl alcohol, and steam distillation had 1.9:1. Furthermore, Nautival [54] stated that organoleptic characteristics are affected by the levels of α - and β-santalol, besides other compounds. Pleasant sandalwood oil extracts are found via SC-CO₂ extraction, hydro, alkaline-hydro, and steam distillation. Furthermore, Nautiyal [55] extracted sandalwood (S. album) oil via SC-CO₂ at 200 bar and 28 °C under two conditions, and the fractionation of the extract was analyzed continuously. Extractions by steam distillation, hydro distillation, Soxhlet extraction were conducted for comparison. The results showed that SC-CO₂ extraction is much more effective in terms of the physical properties of oil than commercial sandalwood oil [55].

Over the last 25 years, about 65,000 chemical structures of the terpenoids and over 7,000 sesquiterpenes (C15) have been reported in previous studies [56]. The EO of the *S. album* tree is composed of the mixture of sesquiterpenes i.e., α -santalol, β -santalol, epi- β -santalol, α -trans-bergamotol, α -bisabolol, lanceol, sesquisabinene hydrate, and farnesol [57]. According to the literature research, α -santalol and β -santalol (Figure 3) which are the main sesquiterpene alcohol compounds found in sandalwood oil are known to indicate biological activities against the skin and prostate cancer and malaria [58]. In the same context, GC analysis of *S. album* oil shows that Z- α -santalol and Z- β -santalol are found with proportions 41–55% and 16–24%, respectively according to the standard [52]; identified some of the sesquiterpenes and monoterpenols, such as α -santalol, β -santalol, α -bergamotol, and cis-lanceol.

According to this chromatographic analysis, santalol levels below these specifications can be related to extraction from undeveloped heartwood, adulteration with synthetic or semi-synthetic substitutes, or substitution with EOs from other species [42].

Mohankumar et al. [59] conducted a study on the heartwood of *S. album* EO concerning antioxidant and stress modulatory efficacy. The traditional steam distillation method is preferred for *S. album* oil extraction and the oil chemical profile identified by the GC–MS technique. *Santalum album* oil has at least 19 main components, accounting for 96.81% of the total content. The main compounds of *S. album* oil followed the order as α -santalol with 41.77% > β -santalol with 18.02% > (Z)- α -trans-bergamotol (8.50%) > (Z)-lanceol (6.57%) > epi- β -santalol (5.78%), cisnuciferol (3.21%) > β -costol (1.41%) > β -santalene (1.24%) > (Z)- β -curcumen-12-ol (1.02%). Besides all components, the pleasant odour of *S. album* oil was contributed by α - and β -santalol.

Subasinghe et al. [60] investigate the Indian sandalwood (*S. album*) EO content and composition in Sri Lanka. Two naturally grown trees heartwoods are studied for comparing the oil properties. The maceration method is applied overnight with deionized water. One of the three the oil yield was measured at 15 cm below ground and found with the highest yield of EO whereas other trees showed a yield varying from 1.46 to 3.35 % w/w.

Another study examines the phytochemical analysis and antibacterial efficiency of extracts of *S. album* in



Figure 3: Structure of α-santalol and β-santalol.

preclinical studies. *In vitro* extracts contain callus, somatic embryo, and seedlings; non-oil-yielding young and oilyielding matured trees are included *in vivo* part. Combined dichloromethane and methanol are used for the 18 h maceration method. Seedlings have the highest amount of sesquiterpenoids with 51.4 mg/g, and the old tree has the least (8.07 mg/g). Monoterpenoids compound content range changes between 3.1 and 4.5 mg/g, except young tree leave extract that has the highest content with 9.5 mg/L [61].

The volatile oil from *S. album* wood and of *Boswellia* sacra Flueck, (syn. *Boswellia carteri* Birdw.) the resin obtained by SC–CO₂ extraction and the effects of extraction conditions on the composition is analyzed in the study of [37]. In general, oxygenated sesquiterpenes dominate the composition of the oil with a 90% ratio and hydrocarbon sesquiterpenes follow these compounds around 5%. According to the results, the best operative conditions is obtained working at 120 bar and 45 °C with the 0.658 g/mL density of CO₂ in the extraction vessels for both matrices.

Another research was conducted on EO composition from roots of *S. album* [62]. Samples were kept in ethyl ether at 48 h for extracting the oil from the root bark. *Santalum album* root heartwood had 10.3% in fresh weight oil yield. Fifty-three different chemical compounds are detected by GC-MS; moreover, β -santalol and α - santalol were included in the ethanolic extract at the highest level with 19.6 and 16%.

In the study of Jones et al. [63]; the yield of the oil from 22 *S. album* trees was evaluated with the use of core sampling at two different heights (30 cm and 100 cm ground level). The results showed that the total concentration of sesquiterpene hydrocarbons is found in a slightly higher proportion in samples. On the other hand, the ratio of α -santalol and β -santalol is lower generally at 100 cm above ground level.

In recent years, procurement of sandalwood resources and their biologically active compounds such as EO or terpenoids have been decreased due to the devastating of the natural stocks and habitats. Therefore, some of the strategies like the heterologous expression, plant cell cultures and plant cell bioreactors have gained prominence to promote the synthesis of sandalwood terpenoids [47, 64].

5.2 Fatty acids

The identification of fatty acid in the oil is generally performed using gas chromatography/mass spectrometry (GC-MS) [65–67]. Zhang et al. [68] examined the 60 compounds from the pericarp-derived volatile oil of *S. album*

with different extraction methods. Colourless EOs are obtained in 2.6 and 5% yield by hydrolyzation and n-hexane extraction and analyzed by GC and GC-MS. Fatty acids, especially palmitic and oleic acids, dominated the total extracted oil with 40–70% depending on the extraction method. *Santalum album* berries proximate analysis, and *in vitro* activities of these compounds have been done by Sri Harsha et al. [69]. Soxhlet method with hexane is used for taking off the oil. The oil that contains a higher amount of oleic acid (45.4%) and palmitic acid (32.5%) is measured 1.5 g/100 g fresh weight. Berries have a very low amount of α -tocopherol when compared to other berry tocopherol content.

5.3 Phenolics and saponins

Santalum album berries phenolic content is found 310 mg gallic acid equivalents (GAE)/100 g fresh weight in methanolic extract of berries. Acidified methanolic extract of *S. album* berries anthocyanin level is measured 0.21% in fresh weight, and the anthocyanin is confirmed as cyanidin 3-glucoside [69]. Various extract of *S. album* has antioxidant activity and a significant role in fighting against free radicals. Kaur et al. [70] reported that the methanolic extract of this strain indicates higher phenolic fractions than other extracts. Besides, cyanidin-3-glucoside is one of the anthocyanin pigments that show antioxidant properties and nutritional potential in *S. album*.

Like *S. album* callus, somatic embryo, and seedlings (*in vitro*); non-oil-yielding young and oil-yielding matured trees (*in vivo*) phenolic results, a similar result is shown in saponin content, *in vivo* extracts with 31.6 and 43.6 mg/g show higher saponin content than *in vitro* extract (9.4 and 17.1 mg/g) [61]. Phytoconstituents and antioxidant activity were analyzed *in vitro* grown callus cultures of *S. album*. The yield of the extract for a dichloromethane–methanol (1:1) solvent mixture was found as 4.3%. The results uncover the abundance of phenolic extracts (18.2 µg). Other major phytoconstituents are found in the extract as terpenoids (16.4 µg/mg), saponins (9.4 µg/mg) and flavan-3-ols (7.0 µg/mg) [71].

Chintamani and Dikshit [72] investigated the antioxidant potential and secondary metabolite of the fruit pulp and the kernels of the *S. album*. As a result of the GC-MS analysis, phenols that have been found in the free form are detected in the acetonitrile extract of fruit kernel and sterol derivatives such as cholest-4-en-3 one compound is recorded mostly in the dichloromethane extract of fruit pulp and. Besides, pyrazine amide and acetamide-2-cyano were obtained as a major constituent of the kernel with the extraction in the methanol and acetonitrile, respectively.

5.4 Phytosterols

β-Sitosterol is found in the *S. album* combined hexane and isopropanol solvent extract (85.35 mg/100 g oil) and *S. album* supercritical CO₂ extract (88.9 mg/100 g oil) at the highest level. Stigmasterol and δ-5-avenasterol amounts are quite higher than other types of chemical compounds [73].

6 Pharmacological activities

6.1 Anticancer activity

Cancer is a population of cell cells with uncontrolled growth and multiplication [74–77]. Natural bioactive compounds help us with anti-inflammatory qualities to fight infections like bacterial, fungal and viral, but also with antioxidant properties with cancer [78–81]. Cytotoxicity is one of the biological activities that characterize sandalwood oil. Mishra et al. [82] in their study showed that new cyclic octapeptide cyclosaplin was cytotoxic against MDA-MB-231 that are human breast cancer cells. Its anticancer activity is based on inducing apoptosis in cells but also on suppression of viability of the cells (Figure 4). Different scientists from the world found that compounds from sandalwood have anticancer activities in many types of skin cancer and leukaemia cells [82-86]. Matsuo and Mimaki [83] found new neolignan and known lignans in sandalwood and this study, they showed that new neolignan was cytotoxic towards HL-60 cells, which are human promyelocytic leukaemia cells. In different work, Matsuo et al. [84] showed that cis- β -santalol and - β -santaldiol were cytotoxic against HL-60 human promyelocytic leukaemia cells by inducing apoptosis in them. According to Santha and Dwivedi [85]; α-santalol from sandalwood oil from Santalum album have anticancer properties, because it can induce apoptosis, have an anti-angiogenic effect and also antioxidant activity on various types of cancer cells.

6.2 Antibacterial, antifungal and antiviral activities

Antibacterial activities are another one that was found among compounds of sandalwood oil due to the content of α/β -santalol and were active against *Salmonella typhimurium* and *Staphylococcus aureus* which are bacteria that cause well known and still threatening diseases throughout the whole world [39, 61, 87]. Epi- β -santalene was found to effective against *S. typhimurium* [88] (Figure 4).

In seeds of Santalum album there is a compound known as santalbic acid, which has antibacterial properties against gram-positive bacteria and antifungal effect on many types of pathogenic fungi [89]. Ochi et al. [90] found that crude organic fractions and sesquiterpenoids from sandalwood oil have antimicrobial activity against Helicobacter pylori which caused peptic ulcers and also can be the cause of gastric cancer. Vadnere et al. [91] purposed to conduct phytochemical analysis and antimicrobial screening of S. album seeds petroleum ether and ethanol extracts. In vitro antimicrobial activity of both extracts was analyzed using a disk diffusion method for Bacillus subtilis, S. aureus, Pseudomonas aeruginosa, Escherichia coli and C. albicans. The outcome of the investigations highlights the potential high efficacy of petroleum ether extract related to santalbic acid, which can function as an antimicrobial agent [91].

Different research showed that derivatives from *S. album* possess significant antifungal properties against species as *Microsporum canis*, *Trichophyton menta-grophytes*, and *Trichophyton rubrum* which is due to the inhibitory effect on mitosis [39, 87, 92]. Powers et al. [93] in their studies have shown that the most active, of the 60 EOs obtained from commercial sources against *Aspergillus niger, Candida albicans*, and *Cryptococcus neoformans*, both in terms of antifungal and cytotoxic activity, were the sandalwood species (*S. album, S. paniculatum*), rich in santalols.

Gupta and Chaphalkar [94] found that aqueous root extract of *S. album* has anti-inflammatory and antiviral activities because it inhibits proliferation, production of monocytes marker – CD14 and also inhibition of nitric oxide in their study on immunopharmalogical activities of it against hepatitis B virus surface antigen – HbsAg, and New Castle disease virus. Antiviral properties were also reported by other scientists in the study of *Herpes simplex* viruses type 1 and 2 [95]. The authors showed that antiviral activity is dose-dependent and didn't exist due to virucidal activity but rather because of the effect on the replication. Sandalwood oil has been also shown to be used against warts, skin blemishes, and other viral-induced tumours on the skin [96].

6.3 Antioxidant

Antioxidants are a group of compounds that protect the body from the chemical process called oxidation [97–99]. This process produces free radicals that attack cell membranes, and for this reason natural antioxidants are important for human health [100, 101]. Antioxidant efficacy is also a known property of sandalwood oil and methanolic extracts from the heartwood of *S. album* [102] but Misra and Dey [71] found it *in vitro* in callus extract of Sandalwood tree and their study showed that it is comparable. Also, antioxidative properties were found in anthocyanins pigment cyanidin-3-glucoside [69].

6.4 Other pharmacological activities

Diabetes is a disease that is widespread along with all the world and sandalwood oil is also found effective in



Figure 4: Illustrative scheme with the most representative pharmacological properties of *Santalum* genus and the correlation with bioactive compounds. Symbols: ↑ increase, ↓ decrease.

DE GRUYTER

managing the complications of this disease [103]. Kulkarni et al. [104] found that it has antihyperglycemic and antihyperlipidemic activities in their studies with diabetic rats, because of the antihyperlipidemic properties it can also help with protecting the liver and also with cardiovascular diseases. Sandalwood extract was reported to inhibit the cardiac tissue damage via reduction of lipid peroxidation damage on the doxorubicin induced cardiotoxicity rat model and significant protective effect against induced myocardial infarction in albino rats in a dose-dependent manner [105].

Sedative activities are known as properties of derivatives from sandalwood [106–108]. Sandalwood oil is reported to produce a relaxing effect on the nerves and is used for headaches, insomnia and nervous tensions. Studies carried out by [109] observed that inhalation of sandalwood oil decreased the motility of mice to an extent of 40–78%. Also showed in their studies that a mild sedative effect occurred in female Swiss albino mice after inhaling sandalwood oil.

Okugawa et al. [107] showed an antipsychotic effect *in vitro* and *in vivo* on mice. In addition, α -santalol is a strong inhibitor of both tyrosinase and cholinesterase *in vivo*, and hence there is a great potential of the EO for use in the treatment of Alzheimer's disease [39].

The potential pharmacological property of *S. album* oil in infective skin conditions have been examined during a few clinical against a wide range of skin conditions. The therapeutic potential of *S. album* oil in dermatology is attributed to its antioxidant, anti-inflammatory and antimicrobial properties. Furthermore, *S. album* oil inhibits the hyper-proliferation of keratinocytes, which is problematic in eczema and psoriasis [110]. Dulal et al. [111] reported that sandalwood oil restores and rejuvenates ageing and wrinkled skin. Sandalwood oil has anti-inflammatory activity as well as emollient used in skincare.

All these pharmacological activities show the value of genus *Santalum* (Figure 4). Sandalwood or sandalwood oil can be used in medicine, cosmetology, and aromatherapy. These innovative materials can solve major issues or diseases such as diabetes, cardiovascular problems, infections of different types, cancer, and also help assist to maintain healthy and beautiful skin and a calm mind.

7 Health-promoting effects: clinical studies

Among all the *Santalum* species, the results of several preclinical studies on *S. album* revealed the vast variety of

pharmaceutical properties of this valuable medicinal plant [61, 91, 94, 112]. Although there are promising *in vitro* and *in vivo* research results on *S. album* oil that shows the high potential capacity of *S. album* oil to treat skin cancer, to date there are limited human studies. Although the available information on sandalwood oil toxicity is limited, it is considered safe due its long history of oral use without any reported adverse effects.

Regarding skin safety, S. album oil has a good safety profile in terms of patch testing for contact dermatitis in both irritation and allergy. According to Burdock and Carabin [43]; undiluted S. album oil and 10% S. album oil are non-irritant. In five dermatology reports, some allergic reactions have been reported. Number of 12 out of 3,542 patients (0.34%) were sensitive to a 2% dilution of S. album oil, and in three reports, 69 of 5,595 patients (1.2%) exhibited sensitivity to a 10% dilution [113]. In a subsequent multicenter European study, 3 fragrance markers (FMs) (fragrance mix I, fragrance mix II, and Myroxylon pereirae) have been tested on consecutive patients to determine the frequency of positive patch-test reactions to EOs tested in the baseline. The result revealed that 656 of 48,956 dermatitis patients (1.38%) revealed positive reactions to 10% S. album oil [114].

Skin inflammation and irritation, known as radiodermatitis, are common side effects in radiation therapy for cancer patients [115]. Radio dermatitis is associated with oxidative stress and an increase in cytokines, including interleukin (IL)-1β, IL-6, and IL-8 [116]. In a study conducted by [117]; the effectiveness of a turmeric and sandalwood oil containing proprietary cream [Vicco[®] turmeric cream (VTC); Vicco Laboratories, Parel, India] on radiodermatitis in patients with head and neck cancer undergoing radiotherapy have been assessed. In this nine-week open-label clinical study, the degree of radiodermatitis of 46 cancer patients experiencing radiotherapy, significantly inhibited (24 patients) compared to baby oil (22 patients) applying Vicco[®] cream containing 16% turmeric extract and 0.5% S. album oil [117]. In a similar study, the same product (Vicco[®]) exhibited significantly delayed and moderated on 40 breast cancer patients (20 in each group) radiodermatitis in the sandalwood/ turmeric group compared to the control group [118].

Based on four clinical trial projects on photoallergy testing, nine of 621 patients (1.45%) tested demonstrated positive effect to *S. album* oil at 2% [119–122]. It should be noted that photoallergy to EOs is very rare, and its clinical application was generally not established.

The potential clinical anti-inflammatory action of sandalwood oil was tested in a clinical trial performed in 50 patients with mild to moderate facial acne for 8 weeks. This pilot study of a topical regimen treatment (foaming cleanser, serum, spot treatment, and mask) containing 0.5% salicylic acid and up to 2% *S. album* oil was conducted in teenage and adult subjects with mild to moderate facial acne [123]. For the eight-week treatment period, treatment was well tolerated by nearly all patients (42 of 47 participants (89.4%). Patients experienced an improvement when compared with baseline with notable reductions in lesion counts in patients with more severe or inflamed lesions, using the Global Aesthetic Improvement Scale (GAIS). There is no report of limitation of use of this regimen due to no adverse events [123].

According to the literature, it is assumed that *S. album* oil might have therapeutic benefits to psoriasis patients due to its anti-inflammatory, antiproliferative properties via inducing autophagy and cell death in proliferating keratinocytes [124–126]. A Phase 2 clinical trial results in patients with mild to moderate psoriasis illustrated that the topically applied 10% *S. album* oil serum administered twice a day for 28 days was well tolerated and alleviates mild to moderate psoriasis symptoms [127].

In a pilot study, undiluted *S. album* oil to common warts twice daily for 12 weeks has been applied to ten candidates with the age range from six to adult. The results showed that 10 of the 12 (80%) participants had complete resolution of all treated warts over their hands, feet, legs, or face, with the other two subjects experiencing moderate improvement. There was no report of skin irritation, redness, pain or other adverse symptoms [128].

8 Safety, adverse effects and therapeutic limitations

Due to the chemical composition of the sandalwood, its EO is most popularly used in folk medicine, cosmetics, pharmacy, as well as the food industry. The list of internal and external health problems, in which the oils of *Santalum* plants are used, contains inter alia general weakness, headache and stomach ache, common colds, bronchitis, skin diseases such as infectious sores, ulcers, acne, and rashes, heart ailments, fever, infection of the urinary tract, and inflammation of the mouth [129, 130].

Just as the positive effect of EOs depend on their chemical composition, their safety and side effects result from the main phytochemicals and as well as the synergistic action of compounds that are present in lower concentrations. The main components present in the sandalwood EOs, that should be taken into the consideration regarding the safety are α -santalol, β -santalol, β -santalene, Z- α -trans-bergamotol [43, 131].

In the food industry, natural flavouring substances may be safely used in the products, meeting some criteria: must be used in the appropriate forms, in the minimum quantity required to produce their intended physical or technical effect, and following all the principles of the good manufacturing practice. Following this, the Food and Drug Administration (FDA) recommendation, a wide range of EO (clove, oregano, thyme, nutmeg, basil, mustard, and cinnamon) and components (linalool, thymol, eugenol, carvone, cinnamaldehyde, vanillin, carvacrol, citral) are classified as generally recognized as safe (GRAS) and have been accepted in the application in food products [132].

According to the FDA, *S. album* is an accepted natural flavouring substance and can be used in the food industry in any kind of product, without restrictions.

Although some limitations with the dosage of the *S. album* EOs are recommended. According to the data of Flavor and Extract Manufacturers Association (FEMA) published in the article of Burdock and Carabin [43] the maximum doses of sandalwood oil in alcoholic beverages should not exceed 0.77 ppm; in non-alcoholic beverages 1.96 ppm, hard candy 89.98 ppm, while in the case of baked products the maximum level is 9.72 ppm. Even though it is difficult to determine how much sandalwood EO is consumed with food by humans, National Academy of Sciences (NAS) data are estimated to 0.0074 mg/day or 0.000123 mg/kg/day sandalwood oil for 60 kg individual.

On the other hand, FEMA reported that these values are 0.0058 mg/day and 0.0001 mg/kg/day, while the mean consumption of foods containing the usual amount- PADI (Possible Average Daily Intake) - is estimated to 0.97 mg/ person/day or 0.016 mg/kg/day of sandalwood oil [43]. In general, the information on the safety and adverse effects of the genus *Santalum* is extremely limited.

Even though the EOs are generally considered as safe, toxicological studies showed that some of them may be harmful to human health. Studies have been shown that different chemicals of EOs (menthol, carvone, limonene, citral, cinnamaldehyde, benzaldehyde, as well as methyl anthranilate, geranyl acetate, furfural, and eugneol) taken at high levels showed no carcinogenic effects [133]. Although low concentrations of EOs are usually devoid of mutagenicity and carcinogenicity, some single components or crude EOs may act as carcinogens. For example, estrogen-dependent malignancy can be induced by *Salvia sclarea* L. EOs, while estragole from *Artemisia dracunculus* L. shows carcinogenic potential in rodents. Following, psoralen (bergamia EOs) is photosensitive compound that may induce DNA adduct formation and skin cancer, while methyleugenol (*Laurus nobilis* L.) and D-limonene (citrus EOs) is being known as carcinogenic in rodents [134].

The lethal dose (LD₅₀) of the sandalwood EOs was evaluated for rats (5.58 g/kg body weight) and rabbits (>5 g/kg BW, body weight). The LD₅₀ was also estimated for the major constituent of the EOs, α -santalol and the values for rats were 3.8 g/kg BW, and for rabbits >5 g/kg BW. 3 mL/kg of α -bisabolol showed a reduction in fetal numbers in rats and rabbits, while 1 mL/kg showed no teratogenic effects [113].

Interestingly, studies on the effects of the inhaled sandalwood oil have shown that female Swiss exposed to the oil for 1 h shown 40% decreased motility, and in the blood of these animals, α - and β -santalols were present [43, 135]. Some studies with animal models suggest that sandalwood EO can irritate rabbit skin, but it seems that this oil has no such effect on human skin [113]. The results of studies showed that sandalwood EO is characterized by low sensitization potential. In the work of Paulsen and Andersen [136] of 318 patients responded that 10% of sandalwood EOs gave a positive effect.

At the same time, 2% concentration did not cause any negative effects on any of the respondents [136]. A total of 1.4% of all tested dentists and dental nurses responded to sandalwood oil in the case of the paper published by Kiecswierczyńska and Krecisz [137]. A total of 0.9% of patients (total of 1606 patients) responded to 10% sandalwood oil, at the same time 0.4% of patients responded to 2% concentration [138].

In the study with 641 patients with eczema, sandalwood oil had no response by any of the tested patients [139], while in the case of 422 patients with suspected contact allergy, 2.4% gave the positive response to sandalwood, and 3.1% to cinnamic alcohol [140]. Similar percentage results were obtained in the studies on the photoallergies' caused by sandalwood oil. 2.2% (3 of 138 patients) were positive to sandalwood oil reaction in the study conducted by Fotiades et al. [119] while 2 of 1050 probable photodermatitis patients (0.19%) in the study of Pigatto et al. [141].

In general sandalwood, EOs are recognized as nontoxic in the matter of phototoxicity, but the suggested maximum dose of the *S. album* EO is 2% [113]. The results of the study with 4266 Japanese people with cosmetic dermatitis showed that 57 (1.34%) was positive to 2% α -santalol. This suggests that the concentration of the sandalwood EOs should be lower for people of Japanese origin [113]. In general, the use of sandalwood oil in eczema, psoriasis, radiation dermatitis, and antifungal is reported in the literature, and the EOs is well tolerated with

acceptable safety [142]. What is more, the major constituent of *S. album* EOs, α -santalol is being recognized as a chemopreventive effect with nontoxic side effects against normal cells [143].

9 Conclusions and future perspectives

The review highlighted the bioactive compounds present in the sandalwood and bioactivities of its extract proven by the in vivo, in vitro and clinical trials. The EO components such as α -santalol and β -santalol are considered important for evaluating the commercial value of the sandalwood. These components are responsible for most of the biological activities along with the soothing aroma of Santalum species. Traditional uses of the EO from sandalwood have been proven to be beneficial in treating somatic and other disorders such as common cold, fever, lung infection, and many types of inflammations. Antioxidant, anti-inflammatory, antibacterial, antifungal, antiviral, neuroleptic, antihyperglycemic, antihyperlipidemic, and anticancer activities of Santalum extracts have been recently proved through clinical trials. Recent scientific studies have not shown any adverse effect of consumption of sandalwood EO in in vivo trials. Hence, extracts from sandalwood are presently used in cosmetic products and as a flavouring agent in food items. More detailed studies are needed to decipher the exact molecular mechanism of the sandalwood extracts in improving human health. These molecular studies will also assist in delineating the more precise use of sandalwood extracts for human consumption. Exhaustive clinical studies are also needed to further promote the use of sandalwood ingredients in food and pharma application. Pharmaceutical formulation of the sandalwood extracts is another area that needs the attention of the scientific community to further improve the use of Santalum species in health promotion.

Acknowledgements: MM wants to thank ANID CENTROS BASALES ACE210012.

Author contributions: All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or in all these areas that is, revising or critically reviewing the article; giving final approval of the version to be published; agreeing on the journal to which the article has been submitted; and confirming to be accountable for all aspects of the work. **Research funding:** None declared.

Competing interest: The author declares no conflict of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

Data availability: All supporting data for this manuscript are included in the figures and available from the corresponding authors on reasonable request.

References

- 1. Subasinghe U. Sandalwood research: a global perspective. J Trop For Environ 2013;3:1–8.
- 2. Fox J. Sandalwood: the royal tree. Biologist 2000;47:31.
- 3. Harbaugh DT, Baldwin BG. Phylogeny and biogeography of the sandalwoods (*Santalum*, Santalaceae): repeated dispersals throughout the Pacific. Am J Bot 2007;94:1028–40.
- Harbaugh D. A taxonomic revision of Australian northern sandalwood (*Santalum lanceolatum*, Santalaceae). Aust Syst Bot 2007;20:409–16.
- Harbaugh DT, Oppenheimer HL, Wood KR, Wagner WL. Taxonomic revision of the endangered Hawaiian red-flowered sandalwoods (*Santalum*) and discovery of an ancient hybrid species. Syst Bot 2010;35:827–38.
- Nageswara rao M, Soneji J, Sudarshana P. Santalum. In: Kole C, editor. Wild crop relatives: genomic and breeding resources. Berlin: Springer; 2011.
- 7. Teixeira da silva JA, Kher MM, Soner D, Page T, Zhang X, Nataraj M, et al. Sandalwood: basic biology, tissue culture, and genetic transformation. Planta 2016;243:847–87.
- Srinivasan V, Sivaramakrishnan V, Rangaswamy C, Ananthapadmanabha H, Shanakaranarayana K. Sandal. Dehra Dun, India: ICFRE; 1992.
- 9. Purohit P. Reviving the royal tree *Santalum album* linn.: Santalaceae. J Med Plants Stud 2018;6:273-6.
- James A. Santalum freycinetianum Gaudich. In: Vozzo J, editor. Tropical tree seed manual: part II, species descriptions. Washington, DC: U.S: Department of Agriculture; 2002.
- Mark D, Thomson L, Elevitch C. Santalum ellipticum, S. freycinetianum, S. haleakalae, and S. paniculatum (Hawaiian sandalwood). In: Elevitch C, editor. Species profiles for Pacific Island agroforestry. ver 4.1. Hawaii: Professional Development Program Grant; 2006.
- Wagner W, Herbst D, Sohmer S. Santalum manual of the flowering plants of Hawaii. Honolulu: University of Hawaii Press; 1999.
- 13. Kepler A. Hawaiian heritage plants. Honolulu, Hawaii, USA: University of Hawaii Press; 1998.
- 14. Wood K. Notes on *Santalum involutum* (Santalaceae) Kaua'i, Hawai'i. *Technical Report*. Hawai'i: National Tropical Botanical Garden; 2015.
- 15. George A. Santalaceae. In: Jessop J, editor. Flora of central Australia. Sydney: Reed Books; 1981.
- George A. Santalaceae. In: George A, editor. Flora of Australia 22 Australian. Canberra: Government Publishing Service; 1984.
- 17. Nge FJ, Ranathunge K, Kotula L, Cawthray GR, Lambers H. Strong host specificity of a root hemi-parasite (*Santalum acuminatum*)

limits its local distribution: beggars can be choosers. Plant Soil 2019;437:159–77.

- Barrett D, Fox J. Geographical distribution of santalaceae and botanical characteristics of species in the genus. In: Cjerum L, Fox J, Ehrhart Y, editors. Sandalwood seed nursery and plantation technology. Proceedings of a regional workshop for Pacific Island countries, 1994, noume'a, new caledonia, RAS field document 8. Suva, Fiji: South Pacific Forestry Development Programme; 1995.
- Cherrier J. Sandalwood in French polynesia. In: Mckinnell F, editor. Sandalwood in the Pacific Region: proceedings of a symposium at the XVII Pacific science congress. Honolulu, Hawaii: Australian Center for International Agriculture Research Proceedings; 1991:24–5 pp.
- 20. Whistler A. Polynesian herbal medicine. Kauai, Hawaii, USA: National Tropical Botanical Garden Press; 1992.
- Payne K. Native plant guide: karlkurla bushland park and the goldfields of Western Australia. Kalgoorlie, Western Australia, Australia: Kalgoorlie-Boulder Urban Landcare Group; 2003.
- Herbst D. Endangered and threatened wildlife and plants; determination of endangered status for *Santalum freycinetianum* var. *lanaiense* (Lanai sandalwood or 'iliahi). Fed Regist 1986;51:3182–5.
- Stuessy TF, Marticorena C, Roberto.rodri guez R, Crawford DJ, Mariosilva O. Endemism in the vascular flora of the Juan Fernandez islands. Aliso 1992;13:297–307.
- 24. PlantList T. Available from: http://www.theplantlist.org/ [Accessed 15 Jun 2022].
- Heinrich M, Appendino G, Efferth T, Fürst R, Izzo AA, Kayser O, et al. Best practice in research – overcoming common challenges in phytopharmacological research. J Ethnopharmacol 2020;246:112230.
- 26. Skottsberg C. The geographical distribution of the sandalwoods and its ignificance. In: Proceedings of the fourth Pacific science congress, java, Indonesia; 1930;vol 3:435–40 pp.
- 27. Stemmermann L. Observations on the genus *Santalum* (Santalaceae) in hawai'i. Pac Sci 1980;34:41–54.
- 28. Tuyama T. On Santalum boninense, and the distribution of the species of *Santalum*. J Jpn Bot 1939;15:697–712.
- Skottsberg C. Further notes on Pacific sandalwoods. Meddelanden från Göteborgs Botaniska Trädgard 1930;5: 135–45.
- Fosberg F, Sachet M. Santalum in eastern polynesia. Candollea 1985;40:459-70.
- Kulkarni H. Sandal (*Santalum album* L.) descriptor. In: Srimathi R, Kulkarni H, KR V, editors. Recent advances in research and management of sandal (*Santalum album* L.) in India. New Delhi: Associated Publishing Company; 1995.
- Srimathi R, Kulkarni H, Venkatesan K. Phenotypes of sandal. J Bombay Nat Hist Soc 1983;80:245–6.
- Lim T. Santalum acuminatum. In: Edible medicinal and nonmedicinal plants. Berlin, Germany: Springer Nature; 2013: 785–9 pp.
- Batabyal S. Growth variability of Santalum album L in different edaphic conditions and formulation of protocols for its propagation [Ph.D. thesis]. Bardhaman, India: University of Burdwan; 2015.
- 35. Chintamani K, Dikshit M. Extraction, antioxidant potential and identification of scondary metabolites of whole fruits of

Santalum album Linn by GC-MS. Int J Curr Pharmaceut Res 2015; 7:40-2.

- Kumar ANA, Joshi G, Ram HYM. Sandalwood: history, uses, present status and the future. Curr Sci 2012;103:1408–16.
- 37. Marongiu B, Piras A, Porcedda S, Tuveri E. Extraction of Santalum album and Boswellia carterii Birdw. volatile oil by supercritical carbon dioxide: influence of some process parameters. Flavour Fragrance J 2006;21:718–24.
- Sindhu R, Arora S. Sandalwood oil: phytochemical and pharmacological updates. In: Govil J, editor. Recent progress in medicinal plants. Studium Press LLC; 2013.
- 39. Misra BB, Dey S. Biological activities of East Indian sandalwood tree, *Santalum album*. PeerJ PrePrints 2013;1:e96v1.
- Scartezzini P, Speroni E. Review on some plants of Indian traditional medicine with antioxidant activity. J Ethnopharmacol 2000;71:23–43.
- Banerjee S, Ecavade A, Rao AR. Modulatory influence of sandalwood oil on mouse hepatic glutathione S-transferase activity and acid soluble sulphydryl level. Cancer Lett 1993;68: 105–9.
- Howes MJ, Simmonds MS, Kite GC. Evaluation of the quality of sandalwood essential oils by gas chromatography-mass spectrometry. J Chromatogr A 2004;1028:307–12.
- 43. Burdock GA, Carabin IG. Safety assessment of sandalwood oil (*Santalum album* L.). Food Chem Toxicol 2008;46:421–32.
- Jones CG, Ghisalberti EL, Plummer JA, Barbour EL. Quantitative co-occurrence of sesquiterpenes; a tool for elucidating their biosynthesis in Indian sandalwood, *Santalum album*. Phytochemistry 2006;67:2463–8.
- Braun NA, Meier M, Pickenhagen W. Isolation and chiral GC analysis of β-bisabolols—trace constituents from the essential Oil of *Santalum album* L. (Santalaceae). J Essent Oil Res 2003;15: 63–5.
- 46. Verghese J, Sunny T, Balakrishnan K. (+)-α-santalol and
 (-)-β-santalol(Z) concentration, a new quality determinant of East Indian sandalwood oil. Flavour Fragrance J 1990;5:2236.
- 47. Zhang Y, Yan H, Niu M, Cheng Q, Zhang X, Teixeira da silva JA, et al. Multiple strategies for increasing yields of essential oil and obtaining sandalwood terpenoids by biotechnological methods in sandalwood. Trees 2018;32:17–28.
- Sharifi-Rad J, Quispe C, Herrera-Bravo J, Akram M, Abbaass W, Semwal P, et al. Phytochemical constituents, biological activities, and health-promoting effects of the *Melissa* officinalis. Oxid Med Cell Longev 2021;2021:6584693.
- 49. Taheri Y, Quispe C, Herrera-Bravo J, Sharifi-Rad J, Ezzat SM, Merghany RM, et al. *Urtica dioica*-derived phytochemicals for pharmacological and therapeutic applications. Evid Based Complement Alternat Med 2022;2022:4024331.
- Salehi B, Rescigno A, Dettori T, Calina D, Docea AO, Singh L, et al. Avocado-soybean unsaponifiables: a panoply of potentialities to Be exploited. Biomolecules 2020;10:130.
- Azwanida N. A review on the extraction methods use in medicinal plants, principle, strength and limitation. Med Aromatic Plants 2015;4:1–6.
- Kusuma HS, Mahfud M. Kinetic studies on extraction of essential oil from sandalwood (*Santalum album*) by microwave airhydrodistillation method. Alex Eng J 2018;57:1163-72.
- 53. Tyśkiewicz K, Gieysztor R, Konkol M, Szałas J, Rój E. Essential oils from humulus lupulus scCO- extract by hydrodistillation and

microwave-assisted hydrodistillation. Molecules 2018;23: 2866.

- Nautiyal O. Subcritical carbon dioxide and conventional extraction techniques of Sandalwood oil: an industry project. Sandalwood Research Newsletter 2010;25:5–8.
- Nautiyal OH. Process optimization of sandalwood (Santalum album) oil extraction by subcritical carbon dioxide and conventional techniques. Indian J Chem Technol 2014;21: 290–7.
- Rusdi NA, Goh HH, Sabri S, Ramzi AB, Mohd noor N, Baharum SN. Functional characterisation of new sesquiterpene synthase from the Malaysian herbal plant, polygonum minus. Molecules 2018; 23. https://doi.org/10.3390/molecules23061370.
- 57. Srivastava PL, Daramwar PP, Krithika R, Pandreka A, Shankar SS, Thulasiram HV. Functional characterization of novel sesquiterpene synthases from Indian sandalwood, *Santalum album*. Sci Rep 2015;5:10095.
- Bommareddy A, Rule B, Vanwert AL, Santha S, Dwivedi C.
 α-Santalol, a derivative of sandalwood oil, induces apoptosis in human prostate cancer cells by causing caspase-3 activation. Phytomedicine 2012;19:804–11.
- Mohankumar A, Kalaiselvi D, Levenson C, Shanmugam G, Thiruppathi G, Nivitha S, et al. Antioxidant and stress modulatory efficacy of essential oil extracted from plantationgrown Santalum album L. Ind Crop Prod 2019;140:111623.
- 60. Subasinghe U, Gamage M, Hettiarachchi DS. Essential oil content and composition of Indian sandalwood (*Santalum album*) in Sri Lanka. J For Res 2013;24:127–30.
- 61. Misra BB, Dey S. Comparative phytochemical analysis and antibacterial efficacy of in vitro and in vivo extracts from East Indian sandalwood tree (*Santalum album* L.). Lett Appl Microbiol 2012;55:476–86.
- Xin-Hua Z, Silva JATD, Yong-Xia J, Jian Y, Guo-Hua M. Essential oils composition from roots of *Santalum album* L. J Essen Oil Bear Plant 2012;15:1–6.
- Jones CG, Plummer JA, Barbour EL. Non-destructive sampling of Indian sandalwood (*Santalum album* L.) for oil content and composition. J Essent Oil Res 2007;19:157–64.
- 64. Rani A, Meghana R, Kush A. Squalene production in the cell suspension cultures of Indian sandalwood (*Santalum album* L.) in shake flasks and air lift bioreactor. Plant Cell Tissue Organ Cult 2018;135:155–67.
- Semwal P, Painuli S, Abu-Izneid T, Rauf A, Sharma A, Daştan SD, et al. Diosgenin: an updated pharmacological review and therapeutic perspectives. Oxid Med Cell Longev 2022;2022: 1035441.
- 66. Quetglas-Llabrés MM, Quispe C, Herrera-Bravo J, Catarino MD, Pereira OR, Cardoso SM, et al. Pharmacological properties of bergapten: mechanistic and therapeutic aspects. Oxid Med Cell Longev 2022;2022:8615242.
- 67. Sharifi-Rad J, Quispe C, Bouyahya A, El-Menyiy N, El Omari N, Shahinozzaman M, et al. Ethnobotany, phytochemistry, biological activities, and health-promoting effects of the genus *Bulbophyllum*. Evid Based Complement Alternat Med 2022; 2022:6727609.
- Zhang XH, Da silva JA, Jia YX, Zhao JT, Ma GH. Chemical composition of volatile oils from the pericarps of Indian sandalwood (*Santalum album*) by different extraction methods. Nat Prod Commun 2012;7:93–6.

- 69. Sri Harsha PSC, Khan MI, Prabhakar P, Giridhar P. Cyanidin-3-glucoside, nutritionally important constituents and in vitro antioxidant activities of *Santalum album* L. berries. Food Res Int 2013;50:275–81.
- 70. Kaur N, Sarkar B, Gill I, Kaur S, Mittal S, Dhiman M, et al. Indian herbs and their therapeutic potential against alzheimer's disease and other neurological disorders. In: Farooqui T, Farooqui A, editors. Neuroprotective effects of phytochemicals in neurological disorders. New Jersey, United States: John Wiley & Sons, Ltd; 2017.
- Misra B, Dey S. Phytochemical analyses and evaluation of antioxidant efficacy of in vitro callus extract of East Indian sandalwood tree (*Santalum album* L.). J Pharmacogn Phytochem 2012;1:8–18.
- Chintamani K, Dikshit M. Extraction, identification and antioxidant potential of secondary metabolites of whole fruits of *Santalum album* Linn by GC-MS. Int J Curr Pharmaceut Res 2015; 7:40–2.
- Hettiarachchi DS, Liu YD, Boddy MR, Fox JED, Sunderland VB. Contents of fatty acids, selected lipids and physicochemical properties of western Australian sandalwood seed oil. J Am Oil Chem Soc 2013;90:285–90.
- 74. Shahinozzaman M, Islam M, Basak B, Sultana A, Emran R, Ashrafizadeh M, et al. A review on chemistry, source and therapeutic potential of lambertianic acid. Z Naturforsch C Biosci 2021;76:347–56.
- Ianoşi SL, Batani A, Ilie MA, Tampa M, Georgescu SR, Zurac S, et al. Non-invasive imaging techniques for the in vivo diagnosis of Bowen's disease: three case reports. Oncol Lett 2019;17: 4094–101.
- Mitrut P, Docea AO, Kamal AM, Mitrut R, Calina D, Gofita E, et al. Colorectal cancer and inflammatory bowel disease. London, United Kingdom: IntechOpen; 2016.
- Zlatian OM, Comanescu MV, Rosu AF, Rosu L, Cruce M, Gaman AE, et al. Histochemical and immunohistochemical evidence of tumor heterogeneity in colorectal cancer. Rom J Morphol Embryol 2015;56:175–81.
- 78. Salehi B, Sharifi-Rad J, Cappellini F, Reiner A, Zorzan D, Imran M, et al. The therapeutic potential of anthocyanins: current approaches based on their molecular mechanism of action. Front Pharmacol 2020b;11:20.
- 79. Sharifi-Rad J, Bahukhandi A, Dhyani P, Sati P, Capanoglu E, Docea AO, et al. Therapeutic potential of neoechinulins and their derivatives: an overview of the molecular mechanisms behind pharmacological activities. Front Nutr 2021;8:664197.
- 80. Dhyani P, Quispe C, Sharma E, Bahukhandi A, Sati P, Attri DC, et al. Anticancer potential of alkaloids: a key emphasis to colchicine, vinblastine, vincristine, vindesine, vinorelbine and vincamine. Cancer Cell Int 2022;22:206.
- Sharifi-Rad J, Quispe C, Patra JK, Singh YD, Panda MK, Das G, et al. Paclitaxel: application in modern oncology and nanomedicine-based cancer therapy. Oxid Med Cell Longev 2021;2021:3687700.
- Mishra A, Gauri SS, Mukhopadhyay SK, Chatterjee S, Das SS, Mandal SM, et al. Identification and structural characterization of a new pro-apoptotic cyclic octapeptide cyclosaplin from somatic seedlings of *Santalum album* L. Peptides 2014;54: 148–58.

- Matsuo Y, Mimaki Y. Lignans from Santalum album and their cytotoxic activities. Chem Pharm Bull (Tokyo) 2010;58: 587–90.
- 84. Matsuo Y, Sakagami H, Mimaki Y. A rare type of sesquiterpene and β-santalol derivatives from *Santalum album* and their cytotoxic activities. Chem Pharm Bull (Tokyo) 2014;62:1192–9.
- 85. Santha S, Dwivedi C. Anticancer effects of sandalwood (*Santalum album*). Anticancer Res 2015;35:3137–45.
- Kim TH, Ito H, Hayashi K, Hasegawa T, Machiguchi T, Yoshida T. Aromatic constituents from the heartwood of *Santalum album* L. Chem Pharm Bull 2005;53:641–4.
- Butaud JF, Gaydou V, Bianchini JP, Faure R, Raharivelomanana P. Dihydroxysesquiterpenoids from *Santalum insulare* of French polynesia. Nat Prod Commun 2007;2:239–42.
- 88. Simanjuntak P. Antibacterial assay of sandalwood (*Santalum album* L.) extract. Majalah Farm 2003;14:326–32.
- Jones GP, Rao KS, Tucker DJ, Richardson B, Barnes A, Rivett DE. Antimicrobial activity of santalbic acid from the oil of *Santalum acuminatum* (quandong). Int J Pharmacogn 1995;33:120–3.
- Ochi T, Shibata H, Higuti T, Kodama KH, Kusumi T, Takaishi Y. Anti-Helicobacter pylori compounds from *Santalum album*. J Nat Prod 2005;68:819–24.
- 91. Vadnere GP, Usman MR, Lodhi S, Patil V. Phytochemical investigation and *in vitro* antimicrobial screening of *Santalum album* seeds extracts. Int J Pharm Pharmaceut Sci 2017;9: 117–24.
- Kim TH, Hatano T, Okamoto K, Yoshida T, Kanzaki H, Arita M, Ito H. Antifungal and Ichthyotoxic sesquiterpenoids from *Santalum album* heartwood. Molecules 22; 2017. https://doi. org/10.3390/molecules22071139.
- Powers CN, Osier JL, Mcfeeters RL, Brazell CB, Olsen EL, Moriarity DM, et al. Antifungal and cytotoxic activities of sixty commercially-available essential oils. Molecules 2018;23: 1549–62.
- 94. Gupta A, Chaphalkar S. Immunopharmacological screening of aqueous root extract of *Santalum album*. J HerbMed Pharma 2015;5:7–11.
- Benencia F, Courrèges M. Antiviral activity of sandalwood oil against Herpes simplex viruses-1 and -2. Phytomedicine 1999;6: 119–23.
- 96. Haque M, Haque A. Use of α and β -santalols, major constituents of sandalwood oil, in the treatment of warts, skin blemishes and other viral- induced tumors. US Patent, 2002:6406706.
- 97. Salehi B, Prakash Mishra A, Nigam M, Karazhan N, Shukla I, Kiełtyka-Dadasiewicz A, et al. *Ficus* plants: state of the art from a phytochemical, pharmacological, and toxicological perspective. Phytother Res 2021;35:1187–217.
- 98. Sharifi-Rad J, Quispe C, Durazzo A, Lucarini M, Souto EB, Santini A, et al. Resveratrol' biotechnological applications: enlightening its antimicrobial and antioxidant properties. J Herb Med 2022b;32:100550.
- 99. Salehi B, Sharifi-Rad J, Capanoglu E, Adrar N, Catalkaya G, Shaheen S, et al. *Cucurbita* plants: from farm to industry. Appl Sci-Basel 2019;9:21.
- 100. Scheau C, Caruntu C, Badarau IA, Scheau AE, Docea AO, Calina D, et al. Cannabinoids and inflammations of the gut-lungskin barrier. J Personalized Med 2021;11. https://doi.org/10. 3390/jpm11060494.

- 101. Salleh WMNHW. A systematic review of botany, phytochemicals and pharmacological properties of "Hoja santa" (*Piper auritum* Kunth). Z Naturforsch C Biosci 2021;76:93–102.
- 102. Saneja A, Kaushik P, Kaushik D, Kumar S, Kumar D. Antioxidant, analgesic and anti-inflammatory activities of *Santalum album* linn. Planta Med 2009;75:452.
- 103. Quispe C, Herrera-Bravo J, Javed Z, Khan K, Raza S, Gulsunoglu-Konuskan Z, et al. Therapeutic applications of curcumin in diabetes: a review and perspective. BioMed Res Int 2022;2022: 1375892.
- 104. Kulkarni CR, Joglekar MM, Patil SB, Arvindekar AU. Antihyperglycemic and antihyperlipidemic effect of *Santalum album* in streptozotocin induced diabetic rats. Pharm Biol 2012; 50:360–5.
- 105. Khan M, Singh M, Khan M, Ahmad S. Protective effect of Santalum album on doxorubicin induced cardiotoxicity in rats. World J Pharmaceut Res 2014;3:2760–71.
- 106. Kumar R, Anjum N, Tripathi YC. Phytochemistry and pharmacology of *Santalum album* L.: a review. World J Pharmaceut Res 2015;4:1842–76.
- Okugawa H, Ueda R, Matsumoto K, Kawanishi K, Kato A. Effect of α-santalol and β-santalol from sandalwood on the central nervous system in mice. Phytomedicine 1995;2: 119–26.
- Okugawa H, Ueda R, Matsumoto K, Kawanishi K, Kato K. Effects of sesquiterpenoids from "Oriental incenses" on acetic acidinduced writhing and D2 and 5-HT2A receptors in rat brain. Phytomedicine 2000;7:417–22.
- 109. Jackson D, Shiju L, Jebasingh D, Huxley V. Memory enhancement potential of *Santalum album* extracts on albino mice. J Theor Biol 2009;5:151.
- Inouye S, Takahashi M, Abe S. Composition, antifungal and radical scavenging activities of 15 rare essential oils. Inter J Essential Oil Therap 2010;4:1–10.
- Dulal S, Taher M, Sheikh H. Sandalwood oil can be miraculous tackle on skin aging, skin appearance and wrinkle skin – a review. World J Pharm Res 2019;5:51–5.
- 112. Ahmed N, Ali khan M, Mat jais A, Mohtarrudin N, Ranjbar M, Amjad M, et al. Anti-ulcer activity of sandalwood (*Santalum album* L.) stem hydroalcoholic extract in three gastric-ulceration models of wistar rats. Bol Latinoam Caribe Plantas Med Aromat 2013;12:81–91.
- 113. Tisserand R, Young R. Essential oil safety: a guide for health care professionals, 2nd ed. St. Louis: Churchill Livingstone; 2014.
- 114. Warshaw EM, Zug KA, Belsito DV, Fowler JF, JR., Dekoven JG, Sasseville D, et al. Positive patch-test reactions to essential oils in consecutive patients from North America and central europe. Derm 2017;28:246–52.
- 115. Sharifi-Rad J, Kamiloglu S, Yeskaliyeva B, Beyatli A, Alfred MA, Salehi B, et al. Pharmacological activities of psoralidin: a comprehensive review of the molecular mechanisms of action. Front Pharmacol 2020;11:11.
- 116. de sanctis V, Agolli L, Visco V, Monaco F, Muni R, Spagnoli A, et al. Cytokines, fatigue, and cutaneous erythema in early stage breast cancer patients receiving adjuvant radiation therapy. BioMed Res Int 2014:523568. https://doi.org/10.1155/2014/ 523568.

- 117. Palatty PL, Azmidah A, Rao S, Jayachander D, Thilakchand KR, Rai MP, et al. Topical application of a sandal wood oil and turmeric based cream prevents radiodermatitis in head and neck cancer patients undergoing external beam radiotherapy: a pilot study. Br J Radiol 2014;87:20130490.
- 118. Rao S, Hegde SK, Baliga-Rao MP, Lobo J, Palatty PL, George T, Baliga MS. Sandalwood oil and turmeric-based cream prevents ionizing radiation-induced dermatitis in breast cancer patients. Clin Study Med 2017;4:43–51.
- 119. Fotiades J, Soter NA, Lim HW. Results of evaluation of 203 patients for photosensitivity in a 7.3-year period. J Am Acad Dermatol 1995;33:597–602.
- 120. Scalf LA, Davis MD, Rohlinger AL, Connolly SM. Photopatch testing of 182 patients: a 6-year experience at the Mayo clinic. Dermatitis 2009;20:44–52.
- Victor FC, Cohen DE, Soter NA. A 20-year analysis of previous and emerging allergens that elicit photoallergic contact dermatitis. J Am Acad Dermatol 2010;62:605–10.
- 122. Greenspoon J, Ahluwalia R, Juma N, Rosen CF. Allergic and photoallergic contact dermatitis: a 10-year experience. Dermatitis 2013;24:29–32.
- 123. Moy RL, Levenson C, So JJ, Rock JA. Single-center, open-label study of a proprietary topical 0.5% salicylic acid-based treatment regimen containing sandalwood oil in adolescents and adults with mild to moderate acne. J Drugs Dermatol JDD 2012;11:1403–8.
- 124. Itoi-Ochi S, Matsumura S, Terao M, Murota H, Katayama I. Sandalwood oil downregulates skin inflammation through 11β-HSD1 activation in keratinocytes. J Dermatol Sci 2016;84: e134.
- 125. Dickinson SE, Olson ER, Levenson C, Janda J, Rusche JJ, Alberts DS, et al. A novel chemopreventive mechanism for a traditional medicine: East Indian sandalwood oil induces autophagy and cell death in proliferating keratinocytes. Arch Biochem Biophys 2014;558:143–52.
- 126. Sharma M, Levenson C, Bell RH, Anderson SA, Hudson JB, Collins CC, et al. Suppression of lipopolysaccharide-stimulated cytokine/chemokine production in skin cells by sandalwood oils and purified α -santalol and β -santalol. Phytother Res 2014;28: 925–32.
- 127. Sharma M, Levenson C, Clements I, Castella P, Gebauer K, Cox ME. East Indian sandalwood oil (EISO) alleviates inflammatory and proliferative pathologies of psoriasis. Front Pharmacol 2017;8:125.
- 128. Haque M, Coury DL. Topical sandalwood oil for common warts. Clin Pediatr (Phila) 2018;57:93–5.
- 129. Preedy VR. Essential oils in food preservation, flavor and safety. San Diego: Academic Press; 2016.
- 130. Goswami NB, Jagatpati T. White sandal (*Santalum album* L.), a precious medicinal and timber yielding plant: a short review. Plant Archives 2018;18:1048–56.
- Misra BB, Dey S. Evaluation of in vivo anti-hyperglycemic and antioxidant potentials of α-santalol and sandalwood oil. Phytomedicine 2013;20:409–16.
- 132. Es I, Khaneghah AM, Akbariirad H. Global regulation of essential oils. In: Essential oils in food processing. New Jersey, United States: John Wiley & Sons, Ltd; 2017.

- 133. Smith RL, Cohen SM, Doull J, Feron VJ, Goodman JI, Marnett LJ, et al. A procedure for the safety evaluation of natural flavor complexes used as ingredients in food: essential oils. Food Chem Toxicol 2005;43:345–63.
- 134. Kuttan R, Liju VB. Safety evaluation of essential oils. New Jersey, United States: John Wiley & Sons, Ltd; 2017.
- 135. Buchbauer G, Jirovetz L, Jäger W, Plank C, Dietrich H. Fragrance compounds and essential oils with sedative effects upon inhalation. J Pharm Sci 1993;82:660–4.
- 136. Paulsen E, Andersen KE. Colophonium and Compositae mix as markers of fragrance allergy: cross-reactivity between fragrance terpenes, colophonium and compositae plant extracts. Contact Derm 2005;53:285–91.
- 137. Kiec-Swierczyńska M, Krecisz B. Allergic contact dermatitis in dentists and dental nurses. Exog Dermatol 2002;1:27–31.
- Frosch PJ, Johansen JD, Menné T, Pirker C, Rastogi SC, Andersen KE, et al. Further important sensitizers in patients sensitive to fragrances. Contact Dermatitis 2002;47: 279–87.

- 139. Trattner A, David M. Patch testing with fine fragrances: comparison with fragrance mix, balsam of Peru and a fragrance series. Contact Derm 2003;49:287–9.
- 140. An S, Lee AY, Lee CH, Kim DW, Hahm JH, Kim KJ, et al. Fragrance contact dermatitis in Korea: a joint study. Contact Dermatitis 2005;53:320–3.
- 141. Pigatto PD, Legori A, Bigardi AS, Guarrera M, Tosti A, Santucci B, et al. Gruppo Italiano Ricerca Dermatiti da Contatto ed Ambientali Italian Multicenter Study of Allergic Contact Photodermatitis: epidemiological aspects. Am J Contact Derm 1996;7:158–63.
- 142. Jain R, Nair S. Sandalwood oil for the chemoprevention of skin cancer: mechanistic insights, anti-inflammatory, and *in vivo* anticancer potential. Current Pharmacology Reports 2019;5: 345–58.
- 143. Bommareddy A, Brozena S, Steigerwalt J, Landis T, Hughes S, Mabry E, et al. Medicinal properties of alpha-santalol, a naturally occurring constituent of sandalwood oil: review. Nat Prod Res 2019;33:527–43.