

A PRELIMINARY SURVEY OF APOIDEA (ANTHOPHILA) AND THEIR USE OF FLORAL RESOURCES ON THE ISLAND OF MO'OREA, FRENCH POLYNESIA

DANIEL H. REJAS

Dept. of Environmental Science, Policy, and Management, University of California, Berkeley, USA

Abstract

The terrestrial biota of the French Polynesian archipelago presents a unique opportunity for study due to a relatively poor understanding of its biology. Among the terrestrial invertebrates, the Apoidea are one of many taxa with incompletely documented biodiversity. This study investigated the diversity of the bees on the island of Mo'orea, part of the Society Islands in French Polynesia. Across a range of elevations, I collected 239 individual bees and observed the floral visits of an additional 266 bees. The visited floral species were recorded to assess apoid use of floral resources, and vegetation surveys of collection sites were conducted to assess the available floral community. A total of five genera of bees were found on Mo'orea, including two which are recorded for the first time in the Society Islands. This study suggests that introduced species, rather than native species, comprise the bee biota of Mo'orea, with the longest established species seemingly introduced at or around the time of colonization by early Polynesians. With the exception of the genera *Lithurgus* and *Megachile*, bee genera were found to rely predominantly on non-native floral resources. Floral visitation predilection by Mo'orean bees may prove to further the spread of introduced and invasive floral species.

Key words: Bee diversity, Apoidea (Anthophila), floral resources, *Lithurgus scabrosus*, *Xylocopa sonorina*, *Megachile umbripennis*, invasive species, Mo'orea, French Polynesia

Introduction

Plant pollination is an essential ecosystem service that serves as a mechanism for the reproduction of most flowering plants. As such, the presence and distributions of flowering plants are the result of successful pollination (Barthell et al. 2001). It is estimated that approximately 60 to 80 percent of the more than 250,000 species of flowering plants on earth depend on animals, primarily insects, for pollination (Kearns & Inouye 1997, Kremen et al. 2007).

Bees, one of the most abundant of floral pollinators, generally are regarded as beneficial for their role in pollination, although negative consequences may exist,

such as pollination of invasive flora and interference with the pollination of native flora (Waser et al. 1996, Simberloff 1999, Richardson et al. 2000, Goulson 2003). Currently, little is known about these potential effects within the French Polynesian archipelago, though studies have been conducted elsewhere in the Pacific, including in New Zealand (Butz Huryn 1997) and Hawaii (Magnacca 2007). The disharmonic nature of island systems of the Pacific, where many important pollinators are absent, can pose challenges for island flora (Gillespie & Roderick 2002, Schueller 2004).

In French Polynesia, the bee fauna is fairly depauperate, due to the relatively


poor dispersal ability of the majority of bee species (Kuhlmann 2006, Michener 2007) and the relative isolation of the Polynesian archipelagos (Gillespie & Roderick 2002). To date, only three species have been quantitatively studied and listed in the Society Islands: *Lithurgus scabrosus* Smith 1853, *Megachile umbripennis* Smith 1874, and *Apis mellifera* Linnaeus 1758 (Michener 1965, Nishida unpubl.). No scientific surveys of Apoidea have been conducted on Mo'orea. For the most part, the apoid taxa of the Pacific islands in general, and French Polynesia more specifically, are generally poorly documented (Kuhlmann 2006, Michener 2007). Comprehensive surveys of bee diversity and biology have been carried out on the Hawaiian islands (Snelling 2003, Magnacca 2007) and the continental landmasses of New Zealand and Australia (Michener 1965, 2007, Goulson & Hanley 2004), but no more than superficial study of the bee fauna has been conducted in the Society Islands since the Bishop Museum's collections in the late 1930's and 1940's (Kuhlmann 2006, pers. comm.).

This study evaluates the apoid biodiversity of Mo'orea, and investigates the use of floral resources by Mo'orea's bees. These preliminary steps toward discerning the roles of Mo'orea's bees in the ecosystem may serve to further planning and conservation efforts for native flora, as well as have implications for our understanding of dispersal and persistence of floral species on islands.

Materials and Methods

Study sites

A total of six study sites were chosen within a range of elevations and vegetation types on the island of Mo'orea, French Polynesia. Collections and observations were conducted between 09 October 2008 and 10 November 2008. I separated sites into three categories in order to investigate potential elevation-based distributions of the bees of



Site	Site type	Location
Afareaitu	Inland	17°32'48" S 149°48'24" W
Kellum estate	Coast	17°29'24" S 149°49'39" W
Lycée agricole	Inland	17°31'33" S 149°50'21" W
Mount Rotui	Ridge	17°29'46" S 149°50'42" W
Temae	Coast	17°29'50" S 149°45'32" W
Trois Cocos	Ridge	17°32'50" S 149°50'30" W

Figure 01. Topographical map of sample sites, with elevations.

Mo'orea. Site categories included low-elevation coastal areas, mid-elevation inland areas, and higher-elevation ridges (Figure 01). At each site, I marked out a 100 meter by 5 meter plot on either side of the access point to the site (e.g. foot trail, path, etc.) within which all floral and hymenopteran specimens were collected and observed.

Vegetation surveys

At each study site, I surveyed the floral species present at each visit within the established plots. Each site survey consisted of 20 one-square-meter quadrats that were evenly spaced across the two plots (Figure 02a), beginning at the edge of the plot and continuing four meters apart. Within each quadrat I documented all floral resources,

identifying to genus and species when possible (Whistler 1996, Welsh 1998, B. Mishler pers. comm.). Flowers were counted individually, except when part of an inflorescence. For floral species with inflorescences, the average number of actual flowers of several inflorescences was counted and multiplied by the number of inflorescences in the quadrat to ascertain the approximate number of flowers.

Specimen collection and observations

I collected bees with a net during a timed one hour period, which occurred predominantly between the hours of 08:30–14:00 at each site. Bees were caught with a net, and transferred into an ethyl acetate kill-jar. The kill-jar was labeled with the floral species the bee was visiting, if applicable. In order to mitigate the transference of pollen between specimens, the kill-jar was lined with a clean kimwipe. Bees were handled and pinned on the same kimwipe, and forceps were cleaned with ethanol between processing of each specimen. Bees were identified to genus, and species if possible (Michener 2007, V. Gonzalez pers. comm.). All insect specimens were sent to and stored at the Essig Museum of Entomology at the University of California, Berkeley.

Following existing recommended protocol for collection of Apoidea, temperature, wind speed, and sky conditions were recorded, with a baseline temperature for sampling set at 18°C for pan traps and 21°C for netting (McCall & Primack 1992, LeBuhn et al. unpubl.). I set out pan traps for five hours per visit at each site, primarily between the hours of 08:30–13:30. 15 pan traps were placed along one of the 100 meter edges of the plots, starting 5 meters in from one end of the plot and then placed every 6 meters. Pan traps were positioned in alternating colors of white, blue, and florescent yellow.

In addition to netting and pan-trapping specimens, I recorded timed observations of




Figure 02a and 02b: Positioning of survey quadrats in relation to site (not to scale).

bee visits to flowers both before and after the netting period. Observation methods were modified from a standardized protocol (Kearns & Inouye 1993) to facilitate comparison across any future studies. Each observation period was 50 minutes long, during which I observed 5 one-square-meter plots for ten minutes each. I recorded a total of 10 sets of observations per site, per visit. Observation plots began 10 meters in from the edge of the plot and then every 20 meters (Figure 02b). During each observation period, I recorded all floral species present in the observational plot, as well as all bee visits to those flowers.

Results

Vegetation surveys

Vegetation surveys yielded a total of 27 floral species, of which 23 were identifiable to at least genus level. I was unable to identify four surveyed floral species, which are labeled Unknown 002, Unknown 004,




Figure 03: Percentage of flowers surveyed, by Family

Unknown 005, and Unknown 017. The species identified to genus comprised 11 plant Families, (Figure 03).

Table 04: Plant community characteristics.

Site	Floral community characteristics [†]			Diversity index [‡]
	Native	Non-native	Unknown	Simpson's index
Afar	0	269	0	4.108
Kell	0	343	31	3.593
Lyc�	0	292	0	4.797
Rotu	12	1	0	1.166
Tema	5	173	6	1.196
Troi	7	289	1	2.537

[†]Plant characters measure in number of flowers.

[‡]Diversity index calculated utilizing all floral species; not a measure of native vs. non-native. The two highest diversity indices are in **bold**.

Over 90% of floral resources surveyed (slightly over 1300 flowers) were from the two plant families Asteraceae (68.6%) and Verbenaceae (22.1%). Floral species known to be native only accounted for 1.8% of floral resources present across all sites, with 95.3% known to be non-native and 2.9% unknown. Simpson's index of diversity $\left[\frac{1}{D}, \text{where } D = \frac{\sum_{i=1}^s n_i(n_i - 1)}{N(N - 1)}\right]$ was calculated to determine a metric to describe floral species abundance in relation to the total species. From these calculations, the inland sites of Afareaitu and Lyc  agricole ranked highest

in diversity, though the flora surveyed at these sites was composed entirely of non-native species (Table 04). For a full list of floral species surveyed, see Appendix A.

The calculation of Simpson's diversity index by site classification type (coastal lowlands, inland mid-elevations, and higher-elevation ridges) yielded the highest index assigned to the inland, mid-elevation sites of Lyc  agricole and Afareaitu. The number of native and non-native floral resources did not show significant statistical differences by site type.

Bee collection and observation

Data on a total of 505 bees was collected from pan traps, timed netting, and observations. The apoid biota recorded included five genera in the two families Apidae and Megachilidae. In Apidae (Apinae), the European honeybee, *Apis mellifera* Linnaeus 1758, was present at all sites sampled. Sampling yielded specimens of *Xylocopa sonorina* Smith 1874, in the family Apidae (Xylocopinae), which was previously unrecorded in the Society Islands. Similarly, the presence of bees in the genus *Ceratina* Latrielle 1802 was recorded for the first time. The two bees in the family Megachilidae, *Lithurgus scabrosus* Smith 1859 and *Megachile umbripennis* Smith 1853, occurred in the lowest numbers, and at the least number of sites (Table 05).

In total, bees utilized 24 different floral

Table 05: Distribution of bees on Mo'orea.

	Afar	Kell	Lyc�	Moun	Tema	Troi
Site type	Inland	Coast	Inland	Ridge	Coast	Ridge
Elevation (m)	63	2	40	405	13	400
Species	Found at site during surveys					
<i>Apis mellifera</i> [254]	Yes	Yes	Yes	Yes	Yes	Yes
<i>Ceratina</i> [223]	Yes	Yes	Yes	No	Yes	Yes
<i>Lithurgus scabrosus</i> [10]	No	No	No	No	Yes	No
<i>Megachile umbripennis</i> [3]	No	No	Yes	Yes	No	No
<i>Xylocopa sonorina</i> [15]	Yes	Yes	Yes	No	Yes	No

Total number of bees of each genus and/or species collected and observed indicated in brackets.

Afar: Afareaitu, Kell: Kellum Estate, Lyc : Lyc  Agricole, Moun: Mount Rotui, Tema: Temae, Troi: Trois Cocos.




Figure 06: Use of native and non-native floral resources by Mo'orea's Apoidea

species, of which 20 were identified to genus and then to species, when possible. Four species that were utilized by bees were not able to be identified, and were labeled as Unknown 002, Unknown 013, Unknown 015, and Unknown 018. The identified plants constituted nine plant families, of which Asteraceae was most heavily utilized by the bee fauna, with 62.6% of all visits occurring on a floral species in Asteraceae.

Similar to the large presence of non-native floral species in the vegetation surveys, non-native species accounted for the bulk of floral visits by the bees of Mo'orea. Of all floral visits recorded by netting from flowers or through observations, a total of 81% of visits occurred on a floral species known to be non-native to the Society Islands, while 14%

of visits occurred on native vegetation (Figure 06). For the genera *Apis* and *Ceratina*, this was found to be statistically significant when examined by a one-way ANOVA performed in JMP version 7.0.2 (*Apis* $p=0.0001$, *Ceratina* $p=0.0072$). For all other genera, sample size did not allow for statistical analysis. Qualitatively, however, use of native and non-native species had significant correlations based on bee genera (Figure 07). Additionally, the overall number of floral visits to non-native species, based on number of floral units present, was found to be statistically significant ($p=0.0382$).

Diagrams of pollinator-plant utilization from the bees collected and observed on Mo'orea indicated high generalization in floral resource use by the bee genera *Apis* and *Ceratina* (Appendix B). The remaining three genera, *X. sonorina*, *L. scabrosus*, and *M. umbripennis*, though collected in relatively low numbers, indicated a stronger specialization in their use of floral resources, with *X. sonorina* collected from two floral species, *M. umbripennis* from two floral species, and *L. scabrosus* solely collected from the native *Hibiscus tiliaceus* (Malvaceae). While low sample size prevented statistical comparison, both *M. umbripennis* and *L. scabrosus* were collected solely on native floral species, while *X. sonorina* was collected solely on non-native floral species.

Simpson's diversity index ranked inland sites highest in terms of diversity of bees by




Figure 07: Percentage of time bee genera collected or observed on native floral species.

site type (Coast 1.509, Inland 3.427, Ridge 1.819). Overall, this correlated with plant diversity, with overall bee diversity highest at sites that were ranked highest in overall plant diversity. Despite low sample size, which prohibited statistical comparison, the general trend existed where greater floral diversity at a survey site corresponded with both greater diversity and higher number of bees at the site.

Bee biology and distribution

Apis mellifera Linnaeus 1758

Distribution: Worldwide. Introduced to the Society Islands (Michener 1965, Kuhlmann 2006, Nishida unpubl.). Collected at all study sites on Mo'orea.

Biology: Eusocial and polylectic. Raised commercially and kept by hobby apiarists on Mo'orea (H. Murphy pers. comm.); feral colonies mostly crevice nesting (Michener 2007).

Lithurgus scabrosus Smith 1853

Distribution: Present in Southeast Asia and through the Pacific Islands to Hawai'i (Michener 1965, Snelling 2003, Kuhlmann 2006). Purportedly a Polynesian introduction to the Polynesian islands (Kuhlmann 2006, Michener 2007).

Biology: Solitary bee that nests primarily in wood of trees, and appears to have narrowly oligolectic relationships with flora in the families Malvaceae and Convolvulaceae (Snelling 2003, Kuhlmann 2006, Michener 2007). Collected solely at Temae from *Hibiscus tiliaceus* (Malvaceae) on Mo'orea.

Megachile umbripennis Smith 1874

Distribution: Present in Southeast Asia and through the Pacific Islands to Hawaii (Michener 1965, 2007, Snelling 2003, Kuhlmann 2006). Suggested to be a Polynesian introduction (Kuhlmann 2006, Michener 2007). Collected at

Lycée Agricole and Mount Rotui on Mo'orea.

Biology: Solitary, wood nesting bee proposed to be narrowly oligolectic on Fabaceae by Kuhlmann (2006). Although one specimen of this species was collected on *Metrosideros collina* (Myrtaceae) at Mount Rotui, specimen was not sampled for *M. collina* pollen.

Xylocopa sonorina Smith 1874

Distribution: Believed to be from the Americas, and species originally described from specimen collected in Hawai'i (Snelling 2003). Present from China, Japan, the Philippines, and through the Pacific islands (Gerling 1983). Not previously described in the Society Islands. Found in coastal and inland elevation sites including Temae, Kellum Estate, and Lycée agricole.

Biology: Solitary to communal nesting bee, making nests in wood (Gerling 1983). Based on foraging behavior on Mo'orea, species may be oligolectic on Verbenaceae and Melastomaceae.

Ceratina Latreille 1802

Distribution: Unlike the other species collected on Mo'orea, I was unable to identify *Ceratina* specimens to species. Two different morphospecies, one or both of which may prove to be *Ceratina dentipes*. Morphospecies 01 larger in size (4-5mm), with two submarginal cells. Facial markings characteristically triangular to hourglass shaped. Morphospecies 02 smaller in size (3.5-4.5mm), with three submarginal cells. Facial markings characteristic rectangular stripe. Specimens of the genus *Ceratina* were collected at all sites except Mount Rotui, though morphospecies varied in collection location (see Appendix B).

Biology: Solitary bees nesting in pithy stems. Nests collected at Afareaitu and Trois Cocos.

A key to the Apoidea of the island of Mo'orea is included in Appendix C.

Discussion

Bee diversity on Mo'orea

The rather depauperate bee biota of Mo'orea includes just five genera, of which four are represented by a single species. Of pollinators, the introduced European honeybee, *A. mellifera*, was the most abundant and ubiquitous of bees, followed closely by those in the genus *Ceratina*. Based on collected and observed specimens both appear fairly cosmopolitan in distribution on the island. Though relatively small numbers of *Xylocopa*, *Lithurgus*, and *Megachile* were collected and observed, certain trends existed in their occurrence. *Xylocopa* occurred in all sites except the high elevation sites, while *Lithurgus* and *Megachile* appeared primarily restricted by the availability of floral resources.

The relatively small number of bee species meshes well with the coupling of the generally poor dispersal ability of bees (Michener 2007) and the effects of the high isolation of islands such as those in French Polynesia (Gillespie & Roderick 2002, Meyer unpubl.). While some of the isolated Polynesian islands have seen adaptive radiation of bee fauna, such as in the native *Hylaeus* (Colletidae) of Hawai'i (Snelling 2003, Magnacca 2007), Mo'orea's bees do not seem to have followed that path. Insular radiations tend toward evolutionary disharmony, skewed by attenuation of solely a few lineages (Gillespie & Roderick 2002). The makeup of Mo'orea's bees, however, is a patchwork of fairly distant and unrelated genera, with the oldest colonizers probably arriving within the last 2000 years.

The diversity of bees was greatest in areas with the highest ranked diversity of plant species. This may be due to the fact that higher floral diversity serves to facilitate pollination through competition

for pollinators, thereby drawing a greater number and diversity of bee species (Ghazoul 2006). Sites that were ranked highest in floral diversity generally had the highest number of floral visitors, though this was most often skewed by the generalist *Apis* and *Ceratina* species. As a generalist pollinator, a site with a diverse floral community may allow for a higher degree of resilience in both bee and flora species, and may equalize the pollen and nectar reward in relation to the cost of travel to other resources (Waser 1996).

Pollinator-plant relationships

The relationship between pollinator and plant carries implications for the persistence and community structure of an ecosystem's flora (Kearns & Inouye 1993, 1997, Ghazoul 2006, Michener 2007). The majority of floral visits by the bees of Mo'orea occurs on non-native plants, and thus may have direct and indirect negative effects on the native flora, either due to competition for pollinators or through synergistic effects among native species. Such synergistic effects, such as invasive "meltdowns," allow facultative exclusion of native species through positive interactions between invasive species. These interactions may include higher levels of pollination or development of floral constancy between or among non-native species (Simberloff & Von Holle 1999, Barthell et al 2001).

While it may be the case that Mo'orea's floral communities originally were predominantly self-pollinated, as is the case in many island systems (Barrett et al. 1996, Schueller 2004), the presence of bees now appears to favor some plant families and species unequally. The generalist bees in high abundance, in the genera *Apis* and *Ceratina*, are predominantly visiting non-native flora, while the Megachilid bees in low abundance, *Lithurgus* and *Megachile*, seem to obligately pollinate native flora. It remains to be seen what the effects of these

visits are on actual plant reproduction and dispersal.

Studies in other ecosystems have documented the ability of *Apis* and generalist bees to act in mutualistic associations with invasive floral species, significantly furthering the spread of invasive species (Howarth 1985, Goulson 2003, Ghazoul 2006). At the same time, however, *Apis* has not been shown to negatively affect the spread of native species (Butz Huryn 1997). In the case of Mo'orea, it may be that, due to the large proportion of non-native floral species present and actively visited by *Apis* and *Ceratina*, these facultative pollination effects and potential mutualistic "meltdowns" may be possible. While visitation of a flower may not necessarily mean pollination occurs, the significantly larger number of non-native species visited may lead to continued or increased dominance of non-native floral species on Mo'orea. Both the abundance and presence of these two bee genera in all types of sites also may indicate that higher elevation refugia for native flora may face further threat from expanding populations of non-native floral species.

The observed use of floral resources by both *Apis* and *Ceratina* greatly outnumbered all other bee species. Patterns of very generalist floral resource use existed – with *Apis* visiting 18 different floral species and *Ceratina* visiting 14 different floral species – and may have important ramifications for plant communities on Mo'orea, especially the aforementioned potential for facultative spreading of non-native flora. The two genera in the family Megachilidae, *Lithurgus* and *Megachile*, appear to be narrowly oligolectic on floral species native to the Polynesian islands (Kuhlmann 2006). As such, they may play an important role in the persistence of native floral species, including *Hibiscus tiliaceus* and *Vigna* species.

Future directions

Present and future interactions between the bee genera and native and non-native floral species carry ramifications for the future structure of the flowering plant communities of Mo'orea. A thorough understanding of the pollination systems present, such as what bees are carrying what pollens, what rates of visitation and pollination success are, and bee population and community structure, must be discerned in order to appropriately conserve Mo'orea's native floral communities and deal with the threat of non-native and invasive flora. Further study is needed to discern the roles of Mo'orea's bee species on these issues, as well as their place within the island ecosystem.

Acknowledgements

I am deeply indebted to the professors, graduate student instructors, and my classmates for their constant support and for providing valuable information and suggestions that greatly shaped this project. Special thanks to the staff of the University of California at Berkeley and the Richard P. Gump South Pacific Research Station who contributed much to the Mo'orea class. Hinano Murphy in particular provided valuable insight into the history of Mo'orea. Genera and species identifications were graciously verified by Dr. Victor Gonzalez (USDA) and from specimens at the Essig Museum of Entomology.

Literature Cited

- Arthur, W. (1995). Wayside plants of the islands: a guide to the lowland flora of the Pacific Islands. University of Hawaii Press. Honolulu HI.
- Barrett, S.C.H., Emerson, B., and Mallet, J. (1996). The reproductive biology and genetics of island plants. *Philosophical Transactions of the Royal Society of London* 351:725-733.

- Barthell, J.F., Randall, J.M., Thorp, R.W., and Wenner, A.W. (2001). Promotion of seed set in yellow star-thistle by honeybees: evidence of an invasive mutualism. *Ecological Applications* 11(6): 1870-1883.
- Butz Huryn, V.M. (1997). Ecological impacts of introduced honey bees. *The Quarterly Review of Biology* 72: 275-297.
- Ghazoul, J. (2006). Floral diversity and the facilitation of pollination. *Journal of Ecology* 94:295-304.
- Gillespie, R.G., and Roderick, G.K. (2002). Arthropods on islands: colonization, speciation, and conservation. *Annual Review of Entomology* 47:595-632.
- Goulson, D. (2003). Effects of introduced bees on native ecosystems. *Annual Review of Ecology and Evolutionary Systems* 34:1-26.
- Howarth, F.G. (1985). Impacts of alien land arthropods and mollusks on native plants and animals in Hawai'i. C.P. Stone & J.M. Scott (eds.). *Hawai'i's Terrestrial Ecosystems*. 149-173. University of HI Press. Honolulu, HI.
- Kearns, C.A. and Inouye, D.W. (1993). *Techniques for pollination biologists*. University Press of Colorado. Niwot, Colorado.
- Kearns, C.A., and Inouye, D.W. (1997). Pollinators, flowering plants, and conservation biology. *BioScience* 47(5):297-307.
- Kremen, C., Williams, N.M., Aizen, M.A., Gemmill-Herren, B., LeBuhn, G., Minckley, R., Packer, L., Potts, S.G., Roulston, T., Steffan-Dewenter, I., Vasquez, D.P., Winfree, R., Adams, L., Crone, E.E., Greenleaf, S., Keitt, T.H., Klein, A., Regetz, J., and Ricketts, T.H. (2007). Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecology Letters* 10:299-314.
- Kuhlmann, M. (2006). Fauna and biogeography of the bees and wasps of the Cook Islands (Hymenoptera aculeata). *Journal of Hymenoptera Research* 15(1):26-37.
- LeBuhn G., Griswold, T., Minckley, R., Droege, S., Roulston, T., Cane, J., Parker, F., Buchmann, S., Tepedino, V., Williams, N., Kremen, C., and Messenger, O. (unpublished). A standardized method for monitoring bee populations – the Bee Inventory (BI) plot. Available at: <http://online.sfsu.edu/~beeplot/>.
- Magnacca, K. (2007). Conservation status of the endemic bees of Hawaii, *Hylaeus* (*Nesoprotopis*) (Hymenoptera: Colletidae). *Pacific Science* 61(2):173-190.
- McCall, C. and Primack, R.B. (1992). Influence of flower characteristics, weather, time of day, and season on insect visitation rates in three plant communities. *American Journal of Botany* 79(4):434-442.
- Meyer, J-Y. (unpublished). Conservation of terrestrial biota in French Polynesia: needs and priorities for research and management.
- Michener, C.D. (1965). A classification of the bees of the Australian and South Pacific regions. *Bulletin of the American museum of Natural History* 130: 1-362.
- Michener, C.D. (2007). *Bees of the world*, 2nd ed. The Johns Hopkins University Press. Baltimore, Maryland.
- Nishida, G.M. (unpublished). French Polynesia ants, bees, and wasps checklist (preliminary), updated 2007. Available: <http://essigdb.berkeley.edu/checklists/>. Accessed: 26 April 2008.
- Richardson, D.M., Allsop, N., D'Antonio, C.M., Milton, S.J., Rejmanek, M. 2000. Plant invasions – the role of mutualisms. *Biological Reviews* 75:65-93.
- Schueller, S.K. 2004. Self-pollination in island and mainland populations of the introduced hummingbird-pollinated plant, *Nicotiana glauca* (Solanaceae). *American Journal of Botany* 91: 672-681.

- Simberloff, D., and Von Holle, B. (1999). Positive interactions of nonindigenous species: invasional meltdown? *Biological Invasions* 1:21-32.
- Snelling, R.R. (2003). Bees of the Hawaiian Islands, exclusive of *Hylaeus* (Nesoprosopis) (Hymenoptera: Apoidea). *Kansas Journal of Entomology* 76(2):342-356.
- Welsh, S.L. (1998). *Flora Societensis: a summary revision of the flowering plants of the Society Islands*. EPS Inc. Orem, Utah.

Appendix A: Floral species surveyed

Plant code	Identification	Sites present	Total floral units [†]
ALP PUR	<i>Alpinia purpurata</i>	Lycée agricole	5
CAL VIA	<i>Calyptocarpus vialis</i>	Kellum estate	13
CLE CHE	<i>Clerodendrum chinense</i>	Kellum estate	7
CRA CRE	<i>Crassocephalum crepedioides</i>	Afareaitu	36
		Kellum estate	9
		Lycée agricole	22
DIS ROT	<i>Dissotis rotundifolia</i>	Afareaitu	1
		Kellum estate	3
ELE MOL	<i>Elephantopus mollis</i>	Afareaitu	23
		Kellum estate	17
		Trois cocos	35
EMI FOS	<i>Emilia fosbergii</i>	Afareaitu	1
		Lycée agricole	28
		Temae	2
		Trois cocos	81
HIB TIL	<i>Hibiscus tiliaceus</i>	Temae	5
		Trois cocos	2
KYL NEM	<i>Kyllinga nemoralis</i>	Kellum estate	2
LAN CAM	<i>Lantana camara</i>	Lycée agricole	3
		Trois cocos	34
MET COL	<i>Metrosideros collina</i>	Mount rotui	12
		Trois cocos	5
MIM PUD	<i>Mimosa pudica</i>	Kellum estate	5
		Lycée agricole	6
MOM CHA	<i>Momordica charantia</i>	Afareaitu	3
		Lycée agricole	21
MOR CIT	<i>Morinda citrifolia</i>	Temae	3
PSE SPI	<i>Pseudoelephantopus spicatus</i>	Afareaitu	78
		Kellum estate	44
PSI SP	<i>Psidium species</i>	Mount rotui	1
RUB ROS	<i>Rubus rosafolium</i>	Kellum estate	6
		Trois cocos	2
SPE SP	<i>Spermacoce species</i>	Kellum estate	3
STA SP	<i>Stachytarpheta species</i>	Afareaitu	29
		Kellum estate	4
		Lycée agricole	13
SYN NOD	<i>Synedrella nodiflora</i>	Afareaitu	4
TRI PRO	<i>Tridax procumbens</i>	Temae	168
VER CIN	<i>Vernonia cineria</i>	Afareaitu	94
		Kellum estate	183
		Lycée agricole	65
		Trois cocos	7
WED SP	<i>Wedelia species</i>	Lycée agricole	1
UNK 002	Unknown 002	Kellum estate	14

UNK 004	Unknown 004	Kellum estate	17
UNK 005	Unknown 005	Trois cocos	1
UNK 017	Unknown 017	Temae	6


† Floral units is a count of the number of flowers or the number of inflorescences for each floral species. If an inflorescence, inflorescence is counted as one floral unit.

Appendix B: Bee genera floral use and collection

Bee species	Sites	Floral species
<i>Apis mellifera</i> (18)	Afareaitu	<i>Clerodendrum chinense</i>
	Kellum Estate	<i>Crassocephalum crepidioides</i>
	Lycée Agricole	<i>Elephantopus mollis</i>
	Mount Rotui	<i>Emilia fosbergii</i>
	Temae	<i>Hibiscus tiliaceus</i>
	Trois Cocos	<i>Lantana camara</i>
		<i>Metrosideros collina</i>
		<i>Momordica charantia</i>
		<i>Morinda citrifolia</i>
		<i>Pseudoelephantopus spicatus</i>
		<i>Psidium</i> species
		<i>Rubus rosifolium</i>
		<i>Spermacoce</i> species
		<i>Stachytarpheta</i> species
		<i>Tridax procumbens</i>
	<i>Vernonia</i> species	
Unknown 002		
<i>Melastoma denticulatum</i>		
<i>Lithurgus scabrosus</i> (1)	Temae	<i>Hibiscus tiliaceus</i>
<i>Megachile umbripennis</i> (2)	Lycée Agricole	<i>Vigna</i> species
	Mount Rotui	<i>Metrosideros collina</i>
<i>Xylocopa sonorina</i> (2)	Afareaitu	<i>Stachytarpheta</i> species
	Kellum Estate	<i>Dissotis rotundifolia</i>
	Lycée Agricole	
	Temae	
<i>Ceratina morphospecies 01</i> (8)	Afareaitu	<i>Crassocephalum crepidioides</i>
	Lycée Agricole	<i>Elephantopus mollis</i>
	Temae	<i>Emilia fosbergii</i>
	Trois Cocos	<i>Lantana camara</i>
		<i>Momordica charantia</i>
		<i>Tridax procumbens</i>
		<i>Vigna</i> species
	Unknown 013	
<i>Ceratina morphospecies 02</i> (10)	Afareaitu	<i>Calyptocarpus vialis</i>
	Kellum Estate	<i>Crassocephalum crepidioides</i>
	Lycée Agricole	<i>Elephantopus mollis</i>
	Temae	<i>Momordica charantia</i>
		<i>Rubus rosifolium</i>
		<i>Spermacoce</i> species
		<i>Synedrella nodiflora</i>
		<i>Tridax procumbens</i>
		<i>Vernonia</i> species
	<i>Dissotis rotundifolia</i>	

Number in parentheses to right of bee species is total number of floral species used.

Pollinator networks with plant families. Plant families with native species on Moorea include Fabaceae, Malvaceae, and Myrtaceae.



Appendix C:

Key to the Apoidea (Anthophila) of Mo'orea, including photos

The bee fauna of Mo'orea is generally depauperate, consisting of only five genera of bees in two families. The below key is for distinguishing among bees solely on the island of Mo'orea, and is intended for use by the amateur entomologist. For scientific keys for identification to genus see Michener 2007 or Agriculture Canada 1993. Species were identified with the help of the UC Berkeley Essig Museum of Entomology and Dr. Victor Gonzalez (USDA).

Key to genus and species

- 1 Small, dark bee, under 8mm in length. Body generally sparsely haired, except for hind tibia. Shiny to dull brown-black. Head with white to yellow facial markings.....(2).

Larger sized bee, over 8mm in length. Hair noticeably present on head, thorax, or abdomen.....(3).
- 2(1) Bee body generally 4-5mm in length. Wing venation with two submarginal cells. Characteristic facial marking in white to yellow, triangular to hourglass shaped.....*Ceratina morphospecies 1*.
[Photos: 01, 02, 03]

Bee body generally 3.5-4.5mm in length, up to 5mm. Wing venation with three submarginal cells. Characteristic facial marking in white to very light yellow, stripe-like with rectangular shape.....*Ceratina morphospecies 2 (C. dentipes?)*.
[Photos: 04, 05]
- 3(1) Large bee, generally over 20mm in length. Generally hairy, with hairless, shiny patch on top of thorax. Densely haired on legs. Females black, males brown to golden color. May be heard buzz-pollinating some plants.....*Xylocopa sonorina* Smith 1874.
[Photos: 06, 07]

Medium sized bee, generally 10-15mm in length. Hairs present either all over body or mainly present on underside of abdomen. Black to yellow-orange in coloration.....(4).
- 4(3) Medium sized bee, generally 10-12mm in length. Typical "bee" coloration, abdomen striped with yellow-orange and black bands. Hairy all over head, thorax, and abdomen. Third tibia with a flattened plate for carrying pollen.....*Apis mellifera* Linnaeus 1758.
[Photos: none]

Medium sized bee, generally 10-14mm in length. Most hair generally restricted to thorax and underside of abdomen. All to mostly black in coloration, or with orange to rust colored hair covering thorax.....(5).
- 5(4) All black bee, usually found visiting flowers of *Hibiscus tiliaceus*. Pollen carried on underside of abdomen on females.....*Lithurgus scabrosus* Smith 1853.
[Photos: 08, 09, 10]

Bee with orange to rust colored hair covering thorax. Head and majority of abdomen all black in color. Pollen carried on underside of abdomen on females. Unlike *A. mellifera*, no striped pattern on abdomen.....*Megachile umbripennis* Smith 1874.
[Photos: 11, 12, 13]

Images

Ceratina morphospecies 1



Photo 01



Photo 02



Photo 03

Ceratina morphospecies 2 (C. dentipes?)



Photo 04




Photo 05

Xylocopa sonorina



Photo 06



Photo 07

Lithurgus scabrosus



Photo 08



Photo 09



Photo 10

Megachile umbripennis



Photo 11



Photo 12



Photo 13

Recommended literature

Agriculture Canada (1993). Hymenoptera of the world: an identification guide to families. Minister of Supplies and Services Canada.

Michener, C.D. (2007). Bees of the world, 2nd ed. The Johns Hopkins University Press. Baltimore, Maryland.