

PAPER • OPEN ACCESS

## Determination spesies flying fishes (*exocoetidae*) in makassar strait

To cite this article: Indrayani *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **441** 012110

View the [article online](#) for updates and enhancements.

You may also like

- [Sustainable flying fish \(\*hirundichthys oxycephalus\*\) fishing with a drift gillnet in Makassar Strait, Indonesia](#)  
Najamuddin, A Assir, M Palo et al.
- [A theoretical analysis of pitch stability during gliding in flying snakes](#)  
Farid Jafari, Shane D Ross, Pavlos P Vlachos et al.
- [Numerical prediction of aerodynamic performance for a flying fish during gliding flight](#)  
Jian Deng, Lingxin Zhang, Zhiyou Liu et al.



**ECS**  
The  
Electrochemical  
Society  
Advancing solid state &  
electrochemical science & technology

**DISCOVER**  
how sustainability  
intersects with  
electrochemistry & solid  
state science research



preservation of flying fish species. This research is expected to be able to represent the diversity of flying fish morphologically in the waters of the Makassar Strait, particularly Majene Regency, West Sulawesi [4]

## 2. Material and methods

This research was conducted in March - April 2019, sampling was collected out in the waters of the Makassar Strait in Majene Regency, and identification of fish samples was carried out in Brawijaya Ichthyologicum Depository, Faculty of Fisheries and Marine Science, Malang. *Exocoetidae* flying fish samples obtained from the location are were documented, and then the recorded samples were identified using the help of a fish identification manual [6]. The documented samples was determined by the type and then written a description of the kind of samples captured, classification of samples from family to species level along with the name of the region. Data collection is carried out by following a fisherman's fishing trip, then during the hauling, a latitude and longitude record is then overlaid into the ArcGIS software.

## 3. Result and discussion

During this research, we found six species of flying fish caught in the Makassar Strait waters. The types of flying fish that has been morphologically identified are as many as four species that represent three genera. The following types of fish were are found in Table 1:

**Table 1.** Species names, Genus and local names

No	Species	Genus	Local Name
1	<i>Cheilopogon spilopterus</i>	<i>Cheilopogon</i>	Tuing-tuing
2	<i>Cheilopogon abei</i>	<i>Cheilopogon</i>	Riri pani
3	<i>Cypselurus poecilopterus</i>	<i>Cypselurus</i>	Tuing-tuing
4	<i>Hirundichthys oxycephalus</i>	<i>Hirundichthys</i>	Torani

### a) *Cheilopogon spilopterus*



**Figure 1.** *Cheilopogon spilopterus*

Maximum standarard length 25 cm. Pelagic fish on the surface close to the coast and neritic. Distribution of this type of fish in the East Indian Ocean and Western Pacific [7].



d) *Hirundichthys oxycephalus***Figure 4.** *Hirundichthys oxycephalus*

The maximum standard length is about 18 cm. Pelagic in nearshore and neritic surface waters. It is a small pelagic fish that is important for Vietnam, Indonesia, and the Philippines. Distributed in the Indian Ocean and the Western Pacific from the Arabian Sea to the southern Solomon Islands, and New South Wales (Australia) [7].

Morphological characters for both genera of *Cheilopogon*, *Cypselurus* and *Hirundichthys* are very similar namely [6] :

1. Lower jaw a little shorter than upper jaw and included beneath the latter; juveniles with a single barbel or not barbelled (*Cypselurus*).
2. Both jaws of equal length, or lower jaw a little longer than upper jaw; juveniles with paired barbels (*Cheilopogon*).
3. Origin of anal fin slightly before, under or not more than 2 rays behind origin of dorsal fin; dorsal fin usually with less, or equal number of rays than anal fin; juveniles not barbelled (*Hirundichthys*).
4. Origin of anal fin 3 rays or more behind origin of dorsal fin; dorsal fin usually with 2 to 5 rays more than anal fin; juveniles barbelled or not barbel.
5. Pelvic fins with a prominent black spot; cross band of pectoral fins usually yellow. (*Cheilopogon abei*).
6. Predorsal scales 24 to 28; pectoral fins with numerous dark spots arranged in regular transverse bands (*Cypselurus poecilopterus*).
7. Predorsal scales 28 to 35; pectoral fins dark brown. (*Cheilopogon spilonotopterus*)
8. Pelvic fins with a prominent black spot; cross band of pectoral fins usually yellow (*Cheilopogon abei*).
9. No palatine teeth; pectoral fin without distinct cross band. (*Hirundichthys oxycephalus*).
10. Origin of anal fin slightly before, under or not more than 2 rays behind origin of dorsal fin; dorsal fin usually with less, or equal number of rays than anal fin; juveniles not barbelled. (*Hirundichthys*).

The distribution of identified flying fish species is obtained from distributed fishing locations as presented in the image map below: Based on the results of the longitude-latitude analysis of the fly fishing area that we collected, then analyzed using Excel and ArcGIS software, the highest-flying fish distribution it was obtained at latitude 177045 ' - 118030' 'BT and 4030'BT - 3005' LS during the study.

**4. Conclusion**

The identification and naming of flying fish groups in Indonesian waters underwent several changes. The discovery of species and these changes are not a severe problem but goes according to time. For this reason, it is recommended to identify flying fish at this time should be followed by DNA analysis for certain species specificity. Also, research on flying fish fisheries in particular to determine the

diversity of species and distribution of their distribution in Indonesia is essential considering the impact of excessive egg exploitation can lead to failure in the recruitment of young fish.

## 5. References

- [1] I B Shakhovskoy and N V Parin 2013 *J. ichthyol.* **53**(2) p 117–118.
- [2] E. A. Lewallen 2012 *tspace.library.utoronto.ca* p 1–197.
- [3] A Syahailatua Febyanti *oseanografi.lipi.go.id*.
- [4] Yahya M Ali 2006 Prediction Potensial Fishing Ground In Makassar Strait *Thesis*.
- [5] Ali S A, M N Nessa, I Djawad and S B A Omar 2004 *Jurnal Ilmu Kelautan dan Perikanan Torani* **3914** 165-172.
- [6] L J V Compagno 1998 *Fao Species Identification Guide For Fishery Purposes Western Central edited by Norwegian Agency for International Development* 3.
- [7] R Parin N V 1999 *Exocoetidae Flyingfishes In K E Carpenter and V H Niem (eds) FAO species identification guide for fishery purposes The living marine resources of the WCP 4 Bony fishes part 2 (Mugilidae to Carangidae)* (Norwegia : FAO) p 2162-2179

## Acknowledgement

Thanks to the Ministry of Research, Technology and Higher Education has provided scholarship and research dissertation doctoral grants in 2019 (No. 1170/K9/KT.03/2018); Dean of the Faculty of Fisheries and Marine Sciences; Director of Postgraduate School, Brawijaya University; who has facilitated, so that this research can do.